



# Bernstein-Type Operators on the Unit Disk

M. J. Recarte<sup>1</sup> · M. E. Marriaga<sup>2</sup> · T. E. Pérez<sup>3</sup>

Received: 25 January 2023 / Revised: 5 May 2023 / Accepted: 10 May 2023  
© The Author(s) 2023

## Abstract

We construct and study sequences of linear operators of Bernstein-type acting on bivariate functions defined on the unit disk. To this end, we study Bernstein-type operators under a domain transformation, we analyze the bivariate Bernstein–Stancu operators, and we introduce Bernstein-type operators on disk quadrants by means of continuously differentiable transformations of the function. We state convergence results for continuous functions and we estimate the rate of convergence. Finally some interesting numerical examples are given, comparing approximations using the shifted Bernstein–Stancu and the Bernstein-type operator on disk quadrants.

**Keywords** Approximation on the ball · Bernstein-type operators

**Mathematics Subject Classification** 32E30 · 41A10 · 41A36

---

Communicated by See Keong Lee.

---

MEM has been supported by the Comunidad de Madrid multiannual agreement with the Universidad Rey Juan Carlos under the Grant Proyectos I+D para Jóvenes Doctores, Ref. M2731, project NETA-MM, and by Ministerio de Ciencia, Innovación y Universidades (MICINN) Grant PGC2018-096504-B-C33 and PID2021-122154NB-I00. TEP and MJR thanks FEDER/Junta de Andalucía under Grant A-FQM-246-UGR20. TEP also thanks MCIN/AEI 10.13039/501100011033 and FEDER funds under Grant PGC2018-094932-B-I00; and IMAG-María de Maeztu Grant CEX2020-001105-M.

---

✉ M. E. Marriaga  
misael.marriaga@urjc.es  
M. J. Recarte  
marlon.recarte@unah.edu.hn  
T. E. Pérez  
tperez@ugr.es

- <sup>1</sup> Departamento de Física, Universidad Nacional Autónoma de Honduras en el valle de Sula, San Pedro Sula, Honduras
- <sup>2</sup> Departamento de Matemática Aplicada, Ciencia e Ingeniería de Materiales y Tecnología Electrónica, Universidad Rey Juan Carlos, Madrid, Spain
- <sup>3</sup> Departamento de Matemática Aplicada, Facultad de Ciencias, Instituto de Matemáticas IMAG, Universidad de Granada, Granada, Spain

## 1 Preliminaries

In 1912, S. Bernstein ([2]) published a constructive proof of the Weierstrass approximation theorem that affirms that every continuous function  $f(x)$  defined on a closed interval can be uniformly approximated by polynomials. For a given function  $f \in C[0, 1]$ , Bernstein constructed a sequence of polynomials (later called Bernstein polynomials) in the form

$$B_n f(x) \equiv (B_n f)(x) = \sum_{k=0}^n f\left(\frac{k}{n}\right) \binom{n}{k} x^k (1-x)^{n-k}, \quad (1.1)$$

for  $0 \leq x \leq 1$ , and  $n \geq 0$ .

Clearly,  $B_n f$  is a polynomial in the variable  $x$  of degree less than or equal to  $n$ , and (1.1) can be seen as a linear operator that transforms functions defined on  $[0, 1]$  to polynomials of degree at most  $n$ .

Hence, in the sequel, we will refer to  $B_n$  as the  $n$ th classical univariate *Bernstein operator*.

If we define

$$p_{n,k}(x) = \binom{n}{k} x^k (1-x)^{n-k}, \quad 0 \leq x \leq 1, \quad 0 \leq k \leq n, \quad (1.2)$$

then the set  $\{p_{n,k}(x) : 0 \leq k \leq n\}$  is a basis of the linear space of polynomials with real coefficients of degree at most  $n$ , that we will denote  $\Pi_n$ , called *Bernstein basis*. Then, the  $n$ th Bernstein polynomial associated with  $f(x)$  is usually written as

$$B_n f(x) = \sum_{k=0}^n f\left(\frac{k}{n}\right) p_{n,k}(x).$$

Among others, classical Bernstein operators satisfy the following properties ([13]):

- They are linear and positive operators acting on the function  $f$  and preserve the constant functions as well as polynomials of degree 1, that is,

$$B_n 1 = 1, \quad B_n x = x, \quad n \geq 0.$$

- If  $f$  is continuous at a point  $x$ , then  $B_n f(x)$  converges to  $f(x)$ , and  $B_n f$  converges uniformly if  $f$  is continuous on the whole interval  $[0, 1]$ . Moreover, the order of approximation is  $\omega_f(n^{-1/2})$ , where  $\omega_f$  denotes the modulus of continuity of  $f$ . Because of this property, Bernstein operators are called *Bernstein Approximants*.
- Bernstein operators satisfy a Voronovskaya-type theorem, that is, if  $f$  is twice differentiable at  $x$ , then  $B_n f(x) - f(x) = \mathcal{O}(1/n)$ .

The Bernstein operators admit a complete system of polynomial eigenfunctions. However, each eigenfunction depends on  $n$  and, thus, is associated with the  $n$ th Bernstein operator  $B_n$ . Another inconvenience of Bernstein operators associated with an adequate function  $f$  is its slow rate of convergence toward  $f$ .

For years, several modifications and extensions of Bernstein operators have been studied. The modifications have been introduced in several directions, and we only recall a few interesting cases and cite some papers. For instance, it is possible to substitute the values of the function on equally spaced points by other mean values such as integrals, as was stated in the pioneering papers of Durrmeyer ([9]) and Derriennic ([6, 7]). In [4], the operator is modified in order to preserve some properties of the original function. Another group of modifications given by the transformation of the function by means a convenient continuous and differentiable functions is analyzed in [5]; and, of course, the extension of the Bernstein operators to the multivariate case. The most common extension of the Bernstein operator is defined on the unit simplex in higher dimensions ([1, 13, 16, 17], among others), since the basic polynomials (1.2) can be easily extended to the simplex.

In this paper, we are interested in finding an extension of the Bernstein operator to approximate functions defined on the unit disk. In this way, we will consider two kinds of modifications: by transformation of the argument of the function to be approximated, and by definition of an adequate basis of functions as (1.2). We present and study two Bernstein-type approximants, and we compare them by means of several examples.

The structure of the paper is as follows. Section 2 is devoted to collecting the properties of univariate Bernstein-type operators that we will need along the paper. In Sect. 3, we recall the method introduced by Stancu ([17]) for obtaining Bernstein-type operators in two variables by the successive application of Bernstein operators in one variable. In Sect. 5 and Sect. 6, we define the *shifted  $n$ th Bernstein–Stancu operator* and the *shifted  $n$ th Bernstein-type operator* and study their respective approximation properties. Section 7 is devoted to describing an extension of certain linear combinations of univariate Bernstein operators that give a better order of approximation. The last section is devoted to analyzing several examples, comparing the approximation results for both Bernstein-type operators on the disk, and the linear combinations introduced in Sect. 7.

## 2 Univariate Bernstein-Type Operators

In this section, we recall the modified univariate Bernstein-type operators that we will need later. We start by shifting the univariate Bernstein operator.

Using the change of variable


$$x = (\beta - \alpha)s + \alpha, \quad \alpha < \beta, \quad 0 \leq s \leq 1, \quad (2.1)$$

the univariate Bernstein basis can be defined on the interval  $[\alpha, \beta]$ . Indeed, if we let

$$\begin{aligned} \tilde{p}_{n,k}(x; [\alpha, \beta]) &= p_{n,k} \left( \frac{x - \alpha}{\beta - \alpha} \right) = \frac{1}{(\beta - \alpha)^n} \binom{n}{k} (x - \alpha)^k (\beta - x)^{n-k}, \\ \alpha &\leq x \leq \beta, \end{aligned}$$

then the set  $\{\tilde{p}_{n,k}(x; [\alpha, \beta]) : n \geq 0, 0 \leq k \leq n, \alpha \leq x \leq \beta\}$  is a basis of  $\Pi_n$  on the interval  $[\alpha, \beta]$  satisfying

**Fig. 1** Bernstein on basis  $[\alpha, \beta]$  for  $n = 5$



$$\begin{aligned} \sum_{k=0}^n \tilde{p}_{n,k}(x; [\alpha, \beta]) &= \sum_{k=0}^n p_{n,k} \left( \frac{x - \alpha}{\beta - \alpha} \right) = \frac{1}{(\beta - \alpha)^n} \sum_{k=0}^n \binom{n}{k} (x - \alpha)^k (\beta - x)^{n-k} \\ &= \frac{1}{(\beta - \alpha)^n} (x - \alpha + \beta - x)^n = 1. \end{aligned}$$

Moreover, since

$$\tilde{p}_{n,k}(x; [\alpha, \beta]) = p_{n,k}(s), \quad 0 \leq s \leq 1, \quad 0 \leq n, \quad 0 \leq k \leq n,$$

we have that Bernstein basis on  $[\alpha, \beta]$  (see Fig. 1) satisfies the following properties:

- $\tilde{p}_{n,k}(x; [\alpha, \beta]) \geq 0$  for  $\alpha \leq x \leq \beta$ ,
- $\tilde{p}_{n,k}(\alpha) = \delta_{0,k}$  and  $\tilde{p}_{n,k}(\beta) = \delta_{k,n}$ , where, as usual,  $\delta_{v,\eta}$  denotes the Kronecker delta,
- $(\beta - \alpha) \tilde{p}'_{n,k}(x; [\alpha, \beta]) = n (\tilde{p}_{n-1,k-1}(x; [\alpha, \beta]) - p_{n-1,k}(x; [\alpha, \beta]))$ ,
- If  $n \neq 0$ , then  $\tilde{p}_{n,k}(x; [\alpha, \beta])$  has a unique local maximum on  $[\alpha, \beta]$  at  $x = (\beta - \alpha) \frac{k}{n} + \alpha$ . This maximum takes the value

$$\tilde{p}_{n,k} \left( (\beta - \alpha) \frac{k}{n} + \alpha; [\alpha, \beta] \right) = p_{n,k} \left( \frac{k}{n} \right) = \binom{n}{k} \frac{k^k}{n^n} (n - k)^{n-k}.$$

For every function  $f$  defined on  $I = [\alpha, \beta]$ , we can define the shifted univariate  $n$ th Bernstein operator as

$$\tilde{B}_n [f(x), I] = \sum_{k=0}^n f \left( (\beta - \alpha) \frac{k}{n} + \alpha \right) \tilde{p}_{n,k}(x; I).$$

Note that  $\tilde{B}_n [f(x), I]$  is a polynomial of degree at most  $n$ . In this way,

$$\tilde{B}_n [f(x), I] = B_n F(s), \quad 0 \leq s \leq 1,$$

where  $F(s) = f((\beta - \alpha)s + \alpha)$  is a function defined on  $[0, 1]$  associated with  $f$ . From this, and since the change of variable (2.1) is linear, it is clear that  $\tilde{B}_n$  has analogous properties to those satisfied by the classical Bernstein operator.

In the sequel, we will use the following Bernstein-type operator studied in [5] and [10]:

$$C_n^\tau f = B_n(f \circ \tau^{-1}) \circ \tau,$$

where  $\tau$  is any function continuously differentiable as many times as necessary, such that  $\tau(0) = 0$ ,  $\tau(1) = 1$ , and  $\tau'(x) > 0$  for  $x \in [0, 1]$ . Throughout this work, it will be sufficient for  $\tau$  to be continuously differentiable.

In [5], the following identities were given:

$$C_n^\tau 1 = 1, \quad C_n^\tau \tau = \tau, \quad C_n^\tau \tau^2 = \left(1 - \frac{1}{n}\right) \tau^2 + \frac{\tau}{n}.$$

We have the following result.

**Proposition 2.1** *Let  $f$  be a continuous function on  $[0, 1]$  and  $\tau$  is any function that is continuously differentiable, such that  $\tau(0) = 0$ ,  $\tau(1) = 1$ , and  $\tau'(x) > 0$  for  $x \in [0, 1]$ . Then,*

$$\lim_{n \rightarrow \infty} C_n^\tau f(x) = f(x).$$

That is,  $C_n^\tau f(x)$  converges uniformly to  $f$  on  $[0, 1]$ .

**Proof** Set  $u = \tau(x)$ . We compute

$$\begin{aligned} C_n^\tau f(x) &= \sum_{k=0}^n f\left(\tau^{-1}\left(\frac{k}{n}\right)\right) p_{n,k}(\tau(x)) \\ &= \sum_{k=0}^n f\left(\tau^{-1}\left(\frac{k}{n}\right)\right) p_{n,k}(u) \\ &= B_n f\left(\tau^{-1}(u)\right). \end{aligned}$$

Since  $B_n f(\tau^{-1}(u)) \rightarrow f(\tau^{-1}(u)) = f(x)$  as  $n \rightarrow +\infty$ , the result follows from taking the limit on both sides of  $C_n^\tau f(x) = B_n f(x)$ .  $\square$

We also introduce the following shifted Bernstein-type operator:

$$\tilde{C}_n^\tau[f(x), [\alpha, \beta]] = \sum_{k=0}^n f \circ \tau^{-1}\left((\beta - \alpha) \frac{k}{n} + \alpha\right) \tilde{p}_{n,k}(\tau(x); [\alpha, \beta]), \quad \alpha \leq x \leq \beta,$$

where  $\tau(x)$  is any function that is continuously differentiable, such that  $\tau(\alpha) = \alpha$ ,  $\tau(\beta) = \beta$ , and  $\tau'(x) > 0$  for  $x \in [\alpha, \beta]$ .

**Proposition 2.2** *Let  $f$  be a continuous function on  $[\alpha, \beta]$  and  $\tau(x)$  is any function that is continuously differentiable, such that  $\tau(\alpha) = \alpha$ ,  $\tau(\beta) = \beta$ , and  $\tau'(x) > 0$  for  $x \in [\alpha, \beta]$ . Then,*

$$\lim_{n \rightarrow \infty} \tilde{C}_n^\tau[f(x), [\alpha, \beta]] = f(x).$$

**Proof** Set  $u = \tau(x)$ . We compute

$$\begin{aligned} \tilde{C}_n^\tau[f(x), [\alpha, \beta]] &= \sum_{k=0}^n f\left(\tau^{-1}\left((\beta - \alpha)\frac{k}{n} + \alpha\right)\right) \tilde{p}_{n,k}(\tau(x); [\alpha, \beta]) \\ &= \sum_{k=0}^n f\left(\tau^{-1}\left((\beta - \alpha)\frac{k}{n} + \alpha\right)\right) \tilde{p}_{n,k}(u; [\alpha, \beta]) \\ &= \tilde{B}_n[f(\tau^{-1}(u)), [\alpha, \beta]]. \end{aligned}$$

Since  $\tilde{B}_n[f(\tau^{-1}(u)), [\alpha, \beta]] \rightarrow f(\tau^{-1}(u)) = f(x)$  as  $n \rightarrow +\infty$ , the result follows from taking the limit on both sides of  $\tilde{C}_n^\tau[f(x), [\alpha, \beta]] = \tilde{B}_n[f(x), [\alpha, \beta]]$ .  $\square$

### 3 Bivariate Bernstein–Stancu operators

In 1963, Stancu [17] studied a method for deducing polynomials of Bernstein type of two variables. This method is based on obtaining an operator in two variables from the successive application of Bernstein operators of one variable.

Let  $\phi_1 \equiv \phi_1(x)$  and  $\phi_2 \equiv \phi_2(x)$  be two continuous functions such that  $\phi_1 < \phi_2$  on  $[0, 1]$ . Let  $\Omega \subseteq \mathbb{R}^2$  be the domain bounded by the curves  $y = \phi_1(x)$ ,  $y = \phi_2(x)$ , and the straight lines  $x = 0$ ,  $x = 1$ . For every function  $f(x, y)$  defined on  $\Omega$ , taking into view

$$y = (\phi_2(x) - \phi_1(x))t + \phi_1(x). \tag{3.1}$$

let us define the function

$$F(x, t) = f(x, (\phi_2(x) - \phi_1(x))t + \phi_1(x)), \tag{3.2}$$

where  $0 \leq t \leq 1$ .

The  $n$ th Bernstein–Stancu operator is defined as

$$\mathcal{B}_n[f(x, y), \Omega] = \sum_{k=0}^n \sum_{j=0}^{n_k} F\left(\frac{k}{n}, \frac{j}{n_k}\right) p_{n,k}(x) p_{n_k,j}(t), \tag{3.3}$$

where each  $n_k$  is a nonnegative integer associated with the  $k$ th node  $x_k = k/n$ , and  $t$  is given by (3.1). Writing (3.3) explicitly, we have

$$\mathcal{B}_n[f(x, y), \Omega] = \sum_{k=0}^n \sum_{j=0}^{n_k} F\left(\frac{k}{n}, \frac{j}{n_k}\right) p_{n,k}(x) p_{n_k,j}\left(\frac{y - \phi_1(x)}{\phi_2(x) - \phi_1(x)}\right),$$

$(x, y) \in \Omega.$

If we denote by  $B_n^{(t)}$  the univariate Bernstein operator acting on the variable  $t$ , then the Bernstein–Stancu operator can be written as

$$\mathcal{B}_n[f(x, y), \Omega] = \sum_{k=0}^n \left[ B_{n_k}^{(t)} F\left(\frac{k}{n}, t\right) \right] p_{n,k}(x).$$

We have the following representation of  $\mathcal{B}_n$  in terms of a matrix determinant.

**Proposition 3.1** *Let  $f(x, y)$  be a function defined on the domain  $\Omega$ , and let  $F$  be a function defined on (3.2). Denote by  $B_n^{(t)}$  the univariate Bernstein operator acting on the variable  $t$ . Then, the  $n$ th Bernstein–Stancu operator is given by the determinant*

$$\mathcal{B}_n[f(x, y), \Omega] = - \left| \begin{array}{ccc|c} 1 & & \circ & B_{n_0}^{(t)} F(0, t) \\ & 1 & & B_{n_1}^{(t)} F(1/n, t) \\ & & \ddots & \vdots \\ \circ & & & 1 \\ \hline & p_{n,0}(x) & p_{n,1}(x) \dots p_{n,n}(x) & B_{n_n}^{(t)} F(1, t) \\ & & & 0 \end{array} \right|.$$

**Remark 3.2** Observe that the step size of the partition of the  $x$ -axis is  $1/n$  and, for a fixed node  $x_k = k/n$ , the step size of the partition of the  $t$ -axis is  $1/n_k$ . Therefore, the step size of the partition of the  $y$ -axis is  $1/m_k$ , where

$$m_k = \frac{n_k}{\phi_2\left(\frac{k}{n}\right) - \phi_1\left(\frac{k}{n}\right)},$$

and, thus,

$$F\left(\frac{k}{n}, \frac{j}{n_k}\right) = f\left(\frac{k}{n}, \frac{j}{m_k} + \phi_1\left(\frac{k}{n}\right)\right).$$

We point out that, in general,  $\mathcal{B}_n[f(x, y), \Omega]$  is not a polynomial. However, it is possible to obtain polynomials by an appropriate choice of  $\phi_1, \phi_2$ , and  $n_k$ . For instance:

- (1) *The Bernstein–Stancu operator on the unit square  $\mathbf{Q} = [0, 1] \times [0, 1]$  (see for instance [13], [17]) is obtained by letting  $\phi_1(x) = 0$  and  $\phi_2(x) = 1$ . Hence, for a function  $f$  defined on  $\mathbf{Q}$ , we get*

$$F\left(\frac{k}{n}, \frac{j}{n_k}\right) = f\left(\frac{k}{n}, \frac{j}{n_k}\right),$$

and

$$\mathcal{B}_n[f(x, y), \mathbf{Q}] = \sum_{k=0}^n \sum_{j=0}^{n_k} f\left(\frac{k}{n}, \frac{j}{n_k}\right) p_{n,k}(x) p_{n_k,j}(y).$$

Note that when  $n_k$  is independent of  $k$  (e.g.,  $n_k = m$  for some positive integer  $m$ ),  $\mathcal{B}_n$  is the tensor product of univariate Bernstein operators on  $\mathbf{Q}$ .

- (2) The Bernstein–Stancu operators can be defined on the simplex  $\mathbf{T}^2 = \{(x, y) \in \mathbb{R}^2 : x, y \geq 0, 1 - x - y \geq 0\}$  (see for instance [1] and [17]). In this case, we set  $\phi_1(x) = 0$ ,  $\phi_2(x) = 1 - x$ , and  $n_k = n - k$ ,  $0 \leq k \leq n$ . In this way,  $m_k = n$  and, since

$$F\left(\frac{k}{n}, \frac{j}{n-k}\right) = f\left(\frac{k}{n}, \frac{j}{n}\right),$$

we have

$$\begin{aligned} \mathcal{B}_n[f(x, y), \mathbf{T}^2] &= \sum_{k=0}^n \sum_{j=0}^{n-k} f\left(\frac{k}{n}, \frac{j}{n}\right) p_{n,k}(x) p_{n-k,j}\left(\frac{y}{1-x}\right) \\ &= \sum_{k=0}^n \sum_{j=0}^{n-k} f\left(\frac{k}{n}, \frac{j}{n}\right) \binom{n}{k \ j} x^k y^j (1-x-y)^{n-k-j}, \\ &\quad (x, y) \in \mathbf{T}^2, \end{aligned}$$

where

$$\binom{n}{k \ j} = \frac{n!}{k! j! (n-k-j)!}, \quad 0 \leq k + j \leq n.$$

In [17], Stancu proved the following convergence result on  $\mathbf{T}^2$ .

**Theorem 3.3** ([17]) *Let  $f$  be a continuous function on  $\mathbf{T}^2$ . Then,  $\mathcal{B}_n[f(x, y), \mathbf{T}^2]$  converges uniformly to  $f(x, y)$  as  $n \rightarrow +\infty$ .*

Stancu only gave a detailed proof of the approximation properties of  $\mathcal{B}_n$  on triangles. In Sect. 5, we consider a slightly general operator and prove the uniform convergence on any bounded domain  $\Omega$ , and we recover Stancu’s result when  $\Omega = \mathbf{T}^2$ .

### 4 Bernstein-Type Operator Under a Domain Transformation

One way to extend the Bernstein operator on the unit square  $\mathbf{Q}$  to another bounded domain  $\Omega \in \mathbb{R}^2$  is through an appropriate transformation or change of variables. In this section, we study several cases.



(1) Let  $\widehat{\mathbf{Q}} = [-1, 1] \times [-1, 1]$ . The operator defined as

$$\widehat{\mathcal{B}}_n[f(x, y), \widehat{\mathbf{Q}}] = \sum_{k=0}^n \sum_{j=0}^{n_k} f\left(\frac{2k-n}{n}, \frac{2j-n_k}{n_k}\right) p_{n,k}\left(\frac{x+1}{2}\right) p_{n_k,j}\left(\frac{y+1}{2}\right),$$

is a Bernstein operator on  $\widehat{\mathbf{Q}}$ . Indeed, for every function  $f$  defined on  $\widehat{\mathbf{Q}}$ , we define the function  $F : \mathbf{Q} \rightarrow \mathbb{R}$  as

$$F(u, v) = f(2u - 1, 2v - 1), \quad (u, v) \in \mathbf{Q}.$$

Then, using the transformation  $x = 2u - 1$  and  $y = 2v - 1$  which maps  $\mathbf{Q}$  into  $\widehat{\mathbf{Q}}$ , we get

$$\begin{aligned} \widehat{\mathcal{B}}_n[f(x, y), \widehat{\mathbf{Q}}] &= \sum_{k=0}^n \sum_{j=0}^{n_k} F\left(\frac{k}{n}, \frac{j}{n_k}\right) p_{n,k}(u) p_{n_k,j}(v) \\ &= \mathcal{B}_n[F(u, v), \mathbf{Q}], \quad (x, y) \in \widehat{\mathbf{Q}}. \end{aligned}$$

(2) An alternative way to obtain the Bernstein–Stancu operator on the simplex  $\mathbf{T}^2$  is by considering the Duffy transformation

$$x = u, \quad y = v(1 - u), \quad (u, v) \in \mathbf{Q},$$

which maps  $\mathbf{Q}$  into  $\mathbf{T}^2$ . Let  $f$  be a function defined on  $\mathbf{T}^2$ . We can define the function  $F : \mathbf{Q} \rightarrow \mathbb{R}$  as

$$F(u, v) = f(u, v(1 - u)), \quad (u, v) \in \mathbf{Q}.$$

Then, the operator

$$\widehat{\mathcal{B}}_n[f(x, y), \mathbf{T}^2] = \sum_{k=0}^n \sum_{j=0}^{n_k} f\left(\frac{k}{n}, \frac{j}{n_k} \left(1 - \frac{k}{n}\right)\right) p_{n,k}(x) p_{n_k,j}\left(\frac{y}{1-x}\right),$$

is a Bernstein-type operator on the simplex since, using the Duffy transformation, we get

$$\widehat{\mathcal{B}}_n[f(x, y), \mathbf{T}^2] = \mathcal{B}_n[F(u, v), \mathbf{Q}], \quad (x, y) \in \mathbf{T}^2.$$

Observe that  $\widehat{\mathcal{B}}_n[f(x, y), \mathbf{T}^2]$  is not a polynomial unless  $n - k - n_k \geq 0$ . We recover the usual Bernstein–Stancu operator on the simplex by setting  $n_k = n - k$ .

(3) Consider the unit ball in  $\mathbb{R}^2$ :

$$\mathbf{B}^2 = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 \leq 1\},$$

and the transformation  $x = 2u - 1, y = (2v - 1) \sqrt{1 - (2u - 1)^2}$  which maps the square  $\mathbf{Q}$  into  $\mathbf{B}^2$ . For every function  $f$  defined on  $\mathbf{B}^2$ , we can define the function  $F : \mathbf{Q} \rightarrow \mathbb{R}^2$  as

$$F(u, v) = f(2u - 1, (2v - 1) \sqrt{1 - (2u - 1)^2}), \quad (u, v) \in \mathbf{Q}.$$

The operator

$$\widehat{\mathcal{B}}_n[f(x, y), \mathbf{B}^2] = \sum_{k=0}^n \sum_{j=0}^{n_k} f\left(\frac{2k - n}{n}, \frac{2j - n_k}{n_k} \frac{2\sqrt{k(n-k)}}{n}\right) p_{n,k}\left(\frac{x+1}{2}\right) p_{n_k,j}\left(\frac{\frac{y}{\sqrt{1-x^2}} + 1}{2}\right),$$

is a Bernstein operator on the unit ball since

$$\widehat{\mathcal{B}}_n[f(x, y), \mathbf{B}^2] = \mathcal{B}_n[F(u, v), \mathbf{Q}], \quad (x, y) \in \mathbf{B}^2.$$

Observe that, in this case,

$$\begin{aligned} & p_{n,k}\left(\frac{x+1}{2}\right) p_{n_k,j}\left(\frac{\frac{y}{\sqrt{1-x^2}} + 1}{2}\right) \\ &= \binom{n}{k} \binom{n_k}{j} \frac{(1+x)^k (1-x)^{n-k} (\sqrt{1-x^2} + y)^j (\sqrt{1-x^2} - y)^{n_k-j}}{2^{n+n_k} (\sqrt{1-x^2})^{n_k}}. \end{aligned}$$

In contrast with the previous two cases, there is no obvious choice of  $n_k$  such that  $\widehat{\mathcal{B}}_n[f(x, y), \mathbf{B}^2]$  is a polynomial. Nevertheless, notice that for  $y = 0$ , we have

$$p_{n,k}\left(\frac{x+1}{2}\right) p_{n_k,j}\left(\frac{1}{2}\right) = \frac{1}{2^{n+n_k}} \binom{n}{k} \binom{n_k}{j} (1+x)^k (1-x)^{n-k},$$

and for  $x = 0$  we have

$$p_{n,k}\left(\frac{1}{2}\right) p_{n_k,j}\left(\frac{y+1}{2}\right) = \frac{1}{2^{n+n_k}} \binom{n}{k} \binom{n_k}{j} (1+y)^j (1-y)^{n_k-j}.$$


Therefore,  $\widehat{\mathcal{B}}_n[f(x, y), \mathbf{B}^2]$  is a polynomial on the  $x$ - and  $y$ -axes for any choice of  $n_k$ .

In Fig. 2, the representation of the mesh in this case for  $n = n_k = 20$  is given.

(4) Let

$$\begin{aligned} B_1 &= \{(x, y) \in \mathbf{B}^2 : x \geq 0, y \geq 0\}, \quad B_2 = \{(x, y) \in \mathbf{B}^2 : x \leq 0, y \geq 0\}, \\ B_3 &= \{(x, y) \in \mathbf{B}^2 : x \leq 0, y \leq 0\}, \quad B_4 = \{(x, y) \in \mathbf{B}^2 : x \geq 0, y \leq 0\}, \end{aligned}$$

**Fig. 2** Mesh corresponding to case (3) for  $n = 20$  and  $n_k = n = 20$  for  $0 \leq k \leq n$



denote the four quadrants of  $\mathbf{B}^2$ , and consider the transformation

$$u = x^2, \quad v = \frac{y^2}{1 - x^2}, \quad (x, y) \in \mathbf{B}^2,$$

which maps each quadrant to  $\mathbf{Q}$ . The corresponding Bernstein operators on the quadrants are:

$$\begin{aligned} \widehat{\mathcal{B}}_n[f(x, y), B_1] &= \sum_{k=0}^n \sum_{j=0}^{n_k} f\left(\sqrt{\frac{k}{n}}, \sqrt{\frac{j(n-k)}{n_k n}}\right) p_{n,k}(x^2) p_{n_k,j}\left(\frac{y^2}{1-x^2}\right), \\ \widehat{\mathcal{B}}_n[f(x, y), B_2] &= \sum_{k=0}^n \sum_{j=0}^{n_k} f\left(-\sqrt{\frac{k}{n}}, \sqrt{\frac{j(n-k)}{n_k n}}\right) p_{n,k}(x^2) p_{n_k,j}\left(\frac{y^2}{1-x^2}\right), \\ \widehat{\mathcal{B}}_n[f(x, y), B_3] &= \sum_{k=0}^n \sum_{j=0}^{n_k} f\left(-\sqrt{\frac{k}{n}}, -\sqrt{\frac{j(n-k)}{n_k n}}\right) p_{n,k}(x^2) p_{n_k,j}\left(\frac{y^2}{1-x^2}\right), \\ \widehat{\mathcal{B}}_n[f(x, y), B_4] &= \sum_{k=0}^n \sum_{j=0}^{n_k} f\left(\sqrt{\frac{k}{n}}, -\sqrt{\frac{j(n-k)}{n_k n}}\right) p_{n,k}(x^2) p_{n_k,j}\left(\frac{y^2}{1-x^2}\right). \end{aligned}$$


Indeed, for every function  $f$  defined on  $\mathbf{B}^2$ , we can define the functions on  $\mathbf{Q}$ :

$$\begin{aligned} F_1(u, v) &= f(\sqrt{u}, \sqrt{v(1-u)}), & F_2(u, v) &= f(-\sqrt{u}, \sqrt{v(1-u)}), \\ F_3(u, v) &= f(-\sqrt{u}, -\sqrt{v(1-u)}), & F_4(u, v) &= f(\sqrt{u}, -\sqrt{v(1-u)}). \end{aligned}$$

Then,

$$\begin{aligned} \widehat{\mathcal{B}}_n[f(x, y), B_1] &= \mathcal{B}_n[F_1(u, v), \mathbf{Q}], & \widehat{\mathcal{B}}_n[f(x, y), B_2] &= \mathcal{B}_n[F_2(u, v), \mathbf{Q}], \\ \widehat{\mathcal{B}}_n[f(x, y), B_3] &= \mathcal{B}_n[F_3(u, v), \mathbf{Q}], & \widehat{\mathcal{B}}_n[f(x, y), B_4] &= \mathcal{B}_n[F_4(u, v), \mathbf{Q}]. \end{aligned}$$

**Fig. 3** Circular mesh after applying the transformation  $(u, v) \mapsto (\sqrt{u}, \sqrt{v(1-u)})$  for  $(u, v) \in \mathbf{Q}$  with  $n = 10$  and  $n_k = n - k$ , for  $0 \leq k \leq n$



If we choose  $n_k = n - k$ , we have that  $\widehat{\mathcal{B}}_n[f(x, y), B_i], i = 1, 2, 3, 4$ , are polynomials of degree  $2n$  since

$$p_{n,k}(x^2) p_{n-k,j} \left( \frac{y^2}{1-x^2} \right) = \binom{n}{k} \binom{n-k}{j} x^{2k} y^{2j} (1-x^2-y^2)^{n-k-j}.$$

In this case, observe that for  $k = 0$ , the mesh corresponding to  $B_1$  and  $B_2$ , and similarly to  $B_3$  and  $B_4$ , coincides on the  $y$ -axis (see Fig. 3). Moreover, for  $j = 0$ , the mesh corresponding to adjacent quadrants coincides on the  $x$ -axis. Therefore, we can define a *piecewise* Bernstein operator on  $\mathbf{B}^2$  as follows:

$$\overline{\mathcal{B}}_n[f(x, y), \mathbf{B}^2] = \begin{cases} \widehat{\mathcal{B}}_n[f(x, y), B_1], & (x, y) \in B_1, \\ \widehat{\mathcal{B}}_n[f(x, y), B_2], & (x, y) \in B_2, \\ \widehat{\mathcal{B}}_n[f(x, y), B_3], & (x, y) \in B_3, \\ \widehat{\mathcal{B}}_n[f(x, y), B_4], & (x, y) \in B_4. \end{cases} \tag{4.1}$$

**Proposition 4.1** *For any function  $f$  on  $\mathbf{B}^2$ ,  $\overline{\mathcal{B}}_n[f(x, y), \mathbf{B}^2]$  is a continuous function on  $\mathbf{B}^2$ .*

**Proof** Clearly,  $\overline{\mathcal{B}}_n[f(x, y), \mathbf{B}^2]$  is continuous on the interior of each quadrant.

For  $x = 0$ ,

$$\begin{aligned} \widehat{\mathcal{B}}_n[f(x, y), B_1] \Big|_{x=0} &= \sum_{k=0}^n \sum_{j=0}^{n-k} f \left( \sqrt{\frac{k}{n}}, \sqrt{\frac{j(n-k)}{n_k n}} \right) p_{n,k}(0) p_{n-k,j}(y^2), \\ &= \sum_{j=0}^{n_0} f \left( 0, \sqrt{\frac{j}{n_0}} \right) p_{n_0,j}(y^2) \\ &= \widehat{\mathcal{B}}_n[f(x, y), B_2] \Big|_{x=0}, \end{aligned}$$

and

$$\begin{aligned} \widehat{\mathcal{B}}_n[f(x, y), B_3] \Big|_{x=0} &= \sum_{k=0}^n \sum_{j=0}^{n_k} f \left( -\sqrt{\frac{k}{n}}, -\sqrt{\frac{j(n-k)}{n_k n}} \right) p_{n,k}(0) p_{n_k,j}(y^2), \\ &= \sum_{j=0}^{n_0} f \left( 0, -\sqrt{\frac{j}{n_0}} \right) p_{n_0,j}(y^2) \\ &= \widehat{\mathcal{B}}_n[f(x, y), B_4] \Big|_{x=0}. \end{aligned}$$

Similarly, for  $y = 0$

$$\widehat{\mathcal{B}}_n[f(x, y), B_1] \Big|_{y=0} = \widehat{\mathcal{B}}_n[f(x, y), B_4] \Big|_{y=0},$$

and

$$\widehat{\mathcal{B}}_n[f(x, y), B_2] \Big|_{y=0} = \widehat{\mathcal{B}}_n[f(x, y), B_3] \Big|_{y=0}.$$

Therefore,  $\widehat{\mathcal{B}}_n[f(x, y), \mathbf{B}^2]$  is continuous on the  $x$ - and  $y$ -axes. □

### 5 Shifted Bernstein–Stancu Operators

Motivated by the examples of Bernstein operators on different domains introduced in the previous section, now we define the *shifted  $n$ th Bernstein–Stancu operator* and study its approximation properties.

Let  $\phi_1$  and  $\phi_2$  be two continuous functions, and let  $I = [a, b]$  be an interval such that  $\phi_1 < \phi_2$  on  $I$ . Let  $\Omega \subset \mathbb{R}^2$  be the domain bounded by the curves  $y = \phi_1(x)$ ,  $y = \phi_2(x)$ , and the straight lines  $x = a$ ,  $x = b$ . Observe that for a fixed  $x \in I$ , the polynomials  $\tilde{p}_{n,k}(y; [\phi_1(x), \phi_2(x)])$ ,  $n \geq 0$ ,  $0 \leq k \leq n$ , constitute a univariate shifted Bernstein basis on the interval  $[\phi_1(x), \phi_2(x)]$ .

For every function  $f(x, y)$  defined on  $\Omega$ , define the function

$$\tilde{F}(u, v; \Omega) = f((b - a)u + a, (\tilde{\phi}_2(u) - \tilde{\phi}_1(u))v + \tilde{\phi}_1(u)), \tag{5.1}$$

where

$$\tilde{\phi}_i(u) = \phi_i((b - a)u + a), \quad i = 1, 2,$$

$0 \leq u \leq 1$ , and  $0 \leq v \leq 1$ .

The shifted  $n$ th Bernstein–Stancu operator is defined as

$$\begin{aligned} \tilde{\mathcal{B}}_n[f(x, y), \Omega] &= \sum_{k=0}^n \sum_{j=0}^{n_k} \tilde{F} \left( \frac{k}{n}, \frac{j}{n_k}; \Omega \right) \tilde{p}_{n,k}(x; I) \tilde{p}_{n_k,j}(y; [\phi_1, \phi_2]), \\ &(x, y) \in \Omega, \end{aligned}$$

where  $n_k = n - k$  or  $n_k = k$  for all  $0 \leq k \leq n$ . Written in terms of the univariate Bernstein basis, we get

$$\tilde{\mathcal{B}}_n[f(x, y), \Omega] = \sum_{k=0}^n \sum_{j=0}^{n_k} \tilde{F}\left(\frac{k}{n}, \frac{j}{n_k}; \Omega\right) p_{n,k}\left(\frac{x-a}{b-a}\right) p_{n_k,j}\left(\frac{y-\phi_1(x)}{\phi_2(x)-\phi_1(x)}\right).$$

The following result plays an important role when studying the convergence of the shifted Bernstein–Stancu operator.

**Lemma 5.1** *Let  $\phi_1$  and  $\phi_2$  be two continuous functions, and let  $I = [a, b]$  be an interval such that  $\phi_1 < \phi_2$  on  $I$ . Let  $\Omega \subset \mathbb{R}^2$  be the domain bounded by the curves  $y = \phi_1(x)$ ,  $y = \phi_2(x)$ , and the straight lines  $x = a$ ,  $x = b$ . Then:*

- (i)  $\tilde{\mathcal{B}}_n[1, \Omega] = 1$ ,
- (ii)  $\tilde{\mathcal{B}}_n[x, \Omega] = x$ ,
- (iii)  $\tilde{\mathcal{B}}_n[y, \Omega] \rightarrow y$  as  $n \rightarrow +\infty$  uniformly on  $[a, b]$ ,
- (iv)  $\tilde{\mathcal{B}}_n[x^2, \Omega] = x^2 + (x - a)(b - x)/n$ ,
- (v)  $\tilde{\mathcal{B}}_n[y^2, \Omega] \rightarrow y^2$  as  $n \rightarrow +\infty$  uniformly on  $[a, b]$ .

**Proof** (i) Obviously  $\tilde{\mathcal{B}}_n[1, \Omega] = 1$ .

(ii) We computerr

$$\begin{aligned} \tilde{\mathcal{B}}_n[x, \Omega] &= \sum_{k=0}^n \sum_{j=0}^{n_k} \tilde{p}_{n,k}(x; I) \tilde{p}_{n_k,j}(y; [\phi_1, \phi_2]) \left( (b-a)\frac{k}{n} + a \right) \\ &= (b-a) \sum_{k=0}^n \tilde{p}_{n,k}(x; I) \frac{k}{n} \left( \sum_{j=0}^{n_k} \tilde{p}_{n_k,j}(y; [\phi_1, \phi_2]) \right) + a \tilde{\mathcal{B}}_n[1, \Omega] \\ &= (b-a) \sum_{k=0}^n \tilde{p}_{n,k}(x; I) \frac{k}{n} + a \\ &= (b-a) \frac{x-a}{b-a} \sum_{k=0}^{n-1} \tilde{p}_{n-1,k}(x; I) + a \\ &= x. \end{aligned}$$

(iii) Observe that

$$\begin{aligned} \sum_{j=0}^{n_k} \tilde{p}_{n_k,j}(y; [\phi_1, \phi_2]) \frac{j}{n_k} &= \sum_{j=1}^{n_k} \binom{n_k}{j} \frac{(y-\phi_1(x))^j (\phi_2(x)-y)^{n_k-j}}{(\phi_2(x)-\phi_1(x))^{n_k}} \frac{j}{n_k} \\ &= \sum_{j=0}^{n_k-1} \binom{n_k-1}{j} \frac{(y-\phi_1(x))^{j+1} (\phi_2(x)-y)^{n_k-1-j}}{(\phi_2(x)-\phi_1(x))^{n_k}} \tag{5.2} \\ &= \frac{y-\phi_1(x)}{\phi_2(x)-\phi_1(x)}. \end{aligned}$$

Therefore, applying the linearity, we get

$$\begin{aligned}
 \tilde{\mathcal{B}}_n[y, \Omega] &= \sum_{k=0}^n \sum_{j=0}^{n_k} \tilde{p}_{n,k}(x; I) \tilde{p}_{n_k,j}(y; [\phi_1, \phi_2]) \\
 &\quad \left[ \left( \tilde{\phi}_2 \left( \frac{k}{n} \right) - \tilde{\phi}_1 \left( \frac{k}{n} \right) \right) \frac{j}{n_k} + \tilde{\phi}_1 \left( \frac{k}{n} \right) \right] \\
 &= \left[ \sum_{k=0}^n \left( \tilde{\phi}_2 \left( \frac{k}{n} \right) - \tilde{\phi}_1 \left( \frac{k}{n} \right) \right) \tilde{p}_{n,k}(x; I) \right] \frac{y - \phi_1(x)}{\phi_2(x) - \phi_1(x)} \\
 &\quad + \sum_{k=0}^n \tilde{\phi}_1 \left( \frac{k}{n} \right) \tilde{p}_{n,k}(x; I) \\
 &= \tilde{B}_n[\phi_2 - \phi_1, I] \frac{y - \phi_1(x)}{\phi_2(x) - \phi_1(x)} + \tilde{B}_n[\phi_1, I],
 \end{aligned}$$

where  $\tilde{B}_n$  denotes the univariate shifted Bernstein operator acting on the variable  $x$ . Since  $\tilde{B}_n$  converges uniformly for a continuous function, we have

$$\begin{aligned}
 \lim_{n \rightarrow +\infty} \tilde{\mathcal{B}}_n[y, \Omega] &= \lim_{n \rightarrow +\infty} \tilde{B}_n[\phi_2 - \phi_1, I] \frac{y - \phi_1(x)}{\phi_2(x) - \phi_1(x)} + \lim_{n \rightarrow +\infty} \tilde{B}_n[\phi_1, I] \\
 &= [\phi_2(x) - \phi_1(x)] \frac{y - \phi_1(x)}{\phi_2(x) - \phi_1(x)} + \phi_1(x) \\
 &= y.
 \end{aligned}$$

(iv) We compute

$$\begin{aligned}
 \tilde{\mathcal{B}}_n[x^2, \Omega] &= \sum_{k=0}^n \sum_{j=0}^{n_k} \tilde{p}_{n,k}(x; I) \tilde{p}_{n_k,j}(y; [\phi_1, \phi_2]) \left( (b-a) \frac{k}{n} + a \right)^2 \\
 &= (b-a)^2 \sum_{k=0}^n \tilde{p}_{n,k}(x; I) \frac{k^2}{n^2} + 2a(b-a) \frac{x-a}{b-a} \sum_{k=0}^{n-1} \tilde{p}_{n-1,k}(x; I) + a^2 \\
 &= (b-a)^2 \left( \frac{n-1}{n} \left( \frac{x-a}{b-a} \right)^2 \sum_{k=0}^{n-2} \tilde{p}_{n-2,k}(x; I) \right. \\
 &\quad \left. + \frac{1}{n} \frac{x-a}{b-a} \sum_{k=0}^{n-1} \tilde{p}_{n-1,k}(x; I) \right) \\
 &\quad + 2a(x-a) + a^2 \\
 &= x^2 + \frac{(x-a)(b-x)}{n}.
 \end{aligned}$$

(v) Finally, if  $f(x, y) = y^2$  in (5.1), then we get

$$\begin{aligned} \tilde{F}\left(\frac{k}{n}, \frac{j}{n_k}; \Omega\right) &= \left[ \left( \tilde{\phi}_2\left(\frac{k}{n}\right) - \tilde{\phi}_1\left(\frac{k}{n}\right) \right) \frac{j}{n_k} + \tilde{\phi}_1\left(\frac{k}{n}\right) \right]^2 \\ &= \left( \tilde{\phi}_2\left(\frac{k}{n}\right) - \tilde{\phi}_1\left(\frac{k}{n}\right) \right)^2 \frac{j^2}{n_k^2} \\ &\quad + 2 \left( \tilde{\phi}_2\left(\frac{k}{n}\right) - \tilde{\phi}_1\left(\frac{k}{n}\right) \right) \tilde{\phi}_1\left(\frac{k}{n}\right) \frac{j}{n_k} + \tilde{\phi}_1\left(\frac{k}{n}\right)^2. \end{aligned}$$

Then,

$$\begin{aligned} \tilde{\mathcal{B}}_n[y^2, \Omega] &= \sum_{k=0}^n \sum_{j=0}^{n_k} \tilde{p}_{n,k}(x; I) \tilde{p}_{n_k,j}(y; [\phi_1, \phi_2]) \left( \tilde{\phi}_2\left(\frac{k}{n}\right) - \tilde{\phi}_1\left(\frac{k}{n}\right) \right)^2 \frac{j^2}{n_k^2} \\ &\quad + 2 \sum_{k=0}^n \sum_{j=0}^{n_k} \tilde{p}_{n,k}(x; I) \tilde{p}_{n_k,j}(y; [\phi_1, \phi_2]) \left( \tilde{\phi}_2\left(\frac{k}{n}\right) - \tilde{\phi}_1\left(\frac{k}{n}\right) \right) \tilde{\phi}_1\left(\frac{k}{n}\right) \frac{j}{n_k} \\ &\quad + \sum_{k=0}^n \sum_{j=0}^{n_k} \tilde{p}_{n,k}(x; I) \tilde{p}_{n_k,j}(y; [\phi_1, \phi_2]) \tilde{\phi}_1\left(\frac{k}{n}\right)^2. \end{aligned}$$

Observe that

$$\begin{aligned} \sum_{j=0}^{n_k} \tilde{p}_{n_k,j}(y; [\phi_1, \phi_2]) \frac{j^2}{n_k^2} &= \frac{n_k - 1}{n_k} \left( \frac{y - \phi_1(x)}{\phi_2(x) - \phi_1(x)} \right)^2 \sum_{j=0}^{n_k-2} \tilde{p}_{n_k-2,j}(y; [\phi_1, \phi_2]) \\ &\quad + \frac{1}{n_k} \frac{y - \phi_1(x)}{\phi_2(x) - \phi_1(x)} \sum_{j=0}^{n_k-1} \tilde{p}_{n_k-1,j}(y; [\phi_1, \phi_2]) \\ &= \frac{n_k - 1}{n_k} \left( \frac{y - \phi_1(x)}{\phi_2(x) - \phi_1(x)} \right)^2 + \frac{1}{n_k} \frac{y - \phi_1(x)}{\phi_2(x) - \phi_1(x)} \\ &= \left( \frac{y - \phi_1(x)}{\phi_2(x) - \phi_1(x)} \right)^2 + \frac{(y - \phi_1(x)) (\phi_2(x) - y)}{n_k (\phi_2(x) - \phi_1(x))^2}. \end{aligned}$$

Together with (5.2), we get

$$\begin{aligned} \tilde{\mathcal{B}}_n[y^2, \Omega] &= \left( \frac{y - \phi_1(x)}{\phi_2(x) - \phi_1(x)} \right)^2 \tilde{B}_n[\phi_2 - \phi_1, I]^2 \\ &\quad + \frac{(y - \phi_1(x)) (\phi_2(x) - y)}{n (\phi_2(x) - \phi_1(x))^2} \sum_{k=0}^n \tilde{p}_{n,k}(x; I) \left( \tilde{\phi}_2\left(\frac{k}{n}\right) - \tilde{\phi}_1\left(\frac{k}{n}\right) \right)^2 \end{aligned}$$



$$\left(\frac{k}{n} - \tilde{\phi}_1\left(\frac{k}{n}\right)\right)^2 \frac{1}{n_k/n} + 2\left(\frac{y - \phi_1(x)}{\phi_2(x) - \phi_1(x)}\right) \tilde{B}_n[(\phi_2 - \phi_1)\phi_1, I] + \tilde{B}_n[\phi_1^2, I].$$

If  $n_k = n - k$ , then

$$\sum_{k=0}^n \tilde{p}_{n,k}(x; I) \left(\tilde{\phi}_2\left(\frac{k}{n}\right) - \tilde{\phi}_1\left(\frac{k}{n}\right)\right)^2 \frac{1}{n_k/n} = \tilde{B}_n\left[\frac{(\phi_2(x) - \phi_1(x))^2}{1 - \frac{x-a}{(b-a)}}, I\right],$$

and if  $n_k = k$ , then

$$\sum_{k=0}^n \tilde{p}_{n,k}(x; I) \left(\tilde{\phi}_2\left(\frac{k}{n}\right) - \tilde{\phi}_1\left(\frac{k}{n}\right)\right)^2 \frac{1}{n_k/n} = \tilde{B}_n\left[\frac{(\phi_2(x) - \phi_1(x))^2}{\frac{x-a}{(b-a)}}, I\right].$$

In either case,  $\tilde{\mathcal{B}}_n[y^2, \Omega] \rightarrow y^2$  as  $n \rightarrow +\infty$ . □

The convergence of the operator is clear from Lemma 5.1 and Volkov’s theorem ([18]).

Now, we study the approximation properties of the shifted Bernstein–Stancu operators.

**Definition 5.2** ([16]) Let  $f$  be a function defined on  $\Omega$ . The modulus of continuity of  $f$  is defined by

$$\omega(\delta_1, \delta_2) = \sup |f(x'', y'') - f(x', y')|,$$

where  $\delta_1, \delta_2 > 0$  are real numbers, whereas  $(x', y')$  and  $(x'', y'')$  are points of  $\Omega$  such that  $|x'' - x'| \leq \delta_1$  and  $|y'' - y'| \leq \delta_2$ .

**Theorem 5.3** Let  $f$  be a continuous function on  $\Omega$ . Then,

$$\lim_{n \rightarrow +\infty} \tilde{\mathcal{B}}_n[f(x, y), \Omega] = f(x, y),$$

uniformly on  $\Omega$ .

**Proof** Let  $\delta_1, \delta_2 > 0$  be real numbers.

Note that on  $\Omega$  we have  $\tilde{\mathcal{B}}_n[1, \Omega] = 1$ ,

$$\tilde{p}_{n,k}(x; I) \tilde{p}_{n_k,j}(y; [\phi_1, \phi_2]) \geq 0, \quad 0 \leq k \leq n, \quad 0 \leq j \leq n_k,$$

and

$$|f(x'', y'') - f(x', y')| \leq \omega(|x'' - x'|, |y'' - y'|) \leq \omega(\delta_1, \delta_2).$$

Taking into account the inequality (see, for instance, [16, 17])

$$\omega(c_1 \delta_1, c_2 \delta_2) \leq (c_1 + c_2 + 1) \omega(\delta_1, \delta_2), \quad c_1, c_2 > 0,$$

we compute

$$\begin{aligned} & \left| f(x, y) - \tilde{F}\left(\frac{k}{n}, \frac{j}{n_k}; \Omega\right) \right| \\ & \leq \omega\left(\left|x - (b-a)\frac{k}{n} - a\right|, \left|y - \left(\tilde{\phi}_2\left(\frac{k}{n}\right) - \tilde{\phi}_1\left(\frac{k}{n}\right)\right)\frac{j}{n_k} - \tilde{\phi}_1\left(\frac{k}{n}\right)\right|\right) \\ & \leq (\lambda_1 + \lambda_2 + 1) \omega(\delta_1, \delta_2), \end{aligned}$$

where

$$\lambda_1 \equiv \lambda_1(x, n, k, \delta_1, a, b) = \frac{1}{\delta_1} \left| x - (b-a)\frac{k}{n} - a \right|,$$

and

$$\lambda_2 \equiv \lambda_2(x, n, k, n_k, \delta_2, \phi_1, \phi_2) = \frac{1}{\delta_2} \left| y - \left(\tilde{\phi}_2\left(\frac{k}{n}\right) - \tilde{\phi}_1\left(\frac{k}{n}\right)\right)\frac{j}{n_k} - \tilde{\phi}_1\left(\frac{k}{n}\right) \right|.$$

Therefore,

$$\begin{aligned} & |f(x, y) - \tilde{\mathcal{B}}_n[f(x, y), \Omega]| \\ & \leq \sum_{k=0}^n \sum_{j=0}^{n_k} \tilde{p}_{n,k}(x; I) \tilde{p}_{n_k,j}(y; [\phi_1, \phi_2]) \left| f(x, y) - \tilde{F}\left(\frac{k}{n}, \frac{j}{n_k}; \Omega\right) \right| \\ & \leq \sum_{k=0}^n \sum_{j=0}^{n_k} \tilde{p}_{n,k}(x; I) \tilde{p}_{n_k,j}(y; [\phi_1, \phi_2]) (\lambda_1 + \lambda_2 + 1) \omega(\delta_1, \delta_2). \end{aligned}$$

We will deal with each term in the last inequality separately.

Since  $\mathcal{B}[1, \Omega] = 1$ ,  $0 \leq \tilde{p}_{n,k}(x; I) \leq 1$ ,  $0 \leq \tilde{p}_{n_k,j}(y; [\phi_1, \phi_2]) \leq 1$ , and  $x \mapsto x^{1/2}$  is a concave function, by Jensen’s inequality, we have

$$\begin{aligned} & \sum_{k=0}^n \sum_{j=0}^{n_k} \tilde{p}_{n,k}(x; I) \tilde{p}_{n_k,j}(y; [\phi_1, \phi_2]) \left| x - (b-a)\frac{k}{n} - a \right| \\ & = \sum_{k=0}^n \sum_{j=0}^{n_k} \tilde{p}_{n,k}(x; I) \tilde{p}_{n_k,j}(y; [\phi_1, \phi_2]) \left[ \left(x - (b-a)\frac{k}{n} - a\right)^2 \right]^{1/2} \\ & \leq \left[ \sum_{k=0}^n \sum_{j=0}^{n_k} \tilde{p}_{n,k}(x; I) \tilde{p}_{n_k,j}(y; [\phi_1, \phi_2]) \left(x - (b-a)\frac{k}{n} - a\right)^2 \right]^{1/2}. \end{aligned}$$

Using (i), (ii), and (iv) in Lemma 5.1, we get

$$\begin{aligned} & \sum_{k=0}^n \sum_{j=0}^{n_k} \tilde{p}_{n,k}(x; I) \tilde{p}_{n_k,j}(y; [\phi_1, \phi_2]) \left| x - (b-a) \frac{k}{n} - a \right| \\ &= \left[ x^2 \tilde{\mathcal{B}}_n[1, \Omega] - 2x \tilde{\mathcal{B}}_n[x, \Omega] + \tilde{\mathcal{B}}_n[x^2, \Omega] \right]^{1/2} \rightarrow 0 \text{ as } n \rightarrow +\infty, \end{aligned}$$

uniformly since  $\tilde{\mathcal{B}}_n[1, \Omega] = 1$ ,  $\tilde{\mathcal{B}}_n[x, \Omega] = x$ , and  $\lim_{n \rightarrow +\infty} \tilde{\mathcal{B}}_n[x^2, \Omega] = x^2$ .

Similarly, from Jensen’s inequality, and using (i), (iii), and (vi) in Lemma 5.1, we get

$$\begin{aligned} & \sum_{k=0}^n \sum_{j=0}^{n_k} \tilde{p}_{n,k}(x; I) \tilde{p}_{n_k,j}(y; \phi_1, \phi_2) \left| y - \left( \tilde{\phi}_2 \left( \frac{k}{n} \right) - \tilde{\phi}_1 \left( \frac{k}{n} \right) \right) \frac{j}{n_k} - \tilde{\phi}_1 \left( \frac{k}{n} \right) \right| \\ & \leq \left[ \sum_{k=0}^n \sum_{j=0}^{n_k} \tilde{p}_{n,k}(x; I) \tilde{p}_{n_k,j}(y; [\phi_1, \phi_2]) \right. \\ & \quad \left. \left( y - \left( \tilde{\phi}_2 \left( \frac{k}{n} \right) - \tilde{\phi}_1 \left( \frac{k}{n} \right) \right) \frac{j}{n_k} - \tilde{\phi}_1 \left( \frac{k}{n} \right) \right)^2 \right]^{1/2} \\ &= \left[ y^2 \tilde{\mathcal{B}}_n[1, \Omega] - 2y \tilde{\mathcal{B}}_n[y, \Omega] + \tilde{\mathcal{B}}_n[y^2, \Omega] \right]^{1/2} \rightarrow 0 \text{ as } n \rightarrow +\infty, \end{aligned}$$

uniformly since  $\tilde{\mathcal{B}}_n[1, \Omega] = 1$ ,  $\lim_{n \rightarrow +\infty} \tilde{\mathcal{B}}_n[y, \Omega] = y$ , and  $\lim_{n \rightarrow +\infty} \tilde{\mathcal{B}}_n[y^2, \Omega] = y^2$ .

Finally, choosing  $\delta_1 = \delta_2 = 1/\sqrt{n}$ , then  $\omega(1/\sqrt{n}, 1/\sqrt{n}) \rightarrow 0$  as  $n \rightarrow +\infty$ , and, thus,  $\tilde{\mathcal{B}}_n[f(x, y), \Omega]$  converges uniformly to  $f(x, y)$  on  $\Omega$ . □

Recall that the univariate shifted Bernstein satisfies the following Voronowskaya type asymptotic formula: Let  $f(x)$  be bounded on the interval  $I$ , and let  $x_0 \in I$  at which  $f''(x_0)$  exists. Then,

$$\tilde{B}_n[f(x), I] \Big|_{x=x_0} - f(x_0) = \mathcal{O} \left( \frac{1}{n} \right). \tag{5.3}$$

Now, we give an analogous result for the Bernstein–Stancu operator.

**Theorem 5.4** *Let  $f(x, y)$  be a bounded function on  $\Omega = \{(x, y) \in \mathbb{R}^2 : a \leq x \leq b, \phi_1(x) \leq y \leq \phi_2(x)\}$ , and let  $(x_0, y_0) \in \Omega$  be a point at which  $f(x, y)$  admits second-order partial derivatives, and  $\phi_i''(x_0)$ ,  $i = 1, 2$ , exist. Then,*

$$\tilde{\mathcal{B}}_n[f(x, y), \Omega] \Big|_{(x,y)=(x_0,y_0)} - f(x_0, y_0) = \mathcal{O} \left( \frac{1}{n} \right).$$

**Proof** Let us write the Taylor expansion of  $f(u, v)$  at the point  $(x_0, y_0)$ :

$$\begin{aligned}
 f(u, v) &= f(x_0, y_0) + (u - x_0) f_x(x_0, y_0) + (v - y_0) f_y(x_0, y_0) \\
 &\quad + \frac{(x - x_0)^2}{2} f_{xx}(x_0, y_0) \\
 &\quad + \frac{(u - x_0)(v - y_0)}{2} (f_{xy}(x_0, y_0) + f_{yx}(x_0, y_0)) + \frac{(v - y_0)^2}{2} f_{yy}(x_0, y_0) \\
 &\quad + \|(u, v) - (x_0, y_0)\|^2 h(u, v),
 \end{aligned}$$

where  $h(u, v)$  is a bounded function such that  $h(u, v) \rightarrow 0$  as  $(u, v) \rightarrow (x_0, y_0)$ . Applying  $\tilde{\mathcal{B}}_n$  to both sides, we get:

$$\begin{aligned}
 \tilde{\mathcal{B}}_n[f(u, v)] &= f(x_0, y_0) + f_x(x_0, y_0) \tilde{\mathcal{B}}_n[u - x_0] + \tilde{\mathcal{B}}_n[v - y_0] f_y(x_0, y_0) \\
 &\quad + \frac{1}{2} f_{xx}(x_0, y_0) \tilde{\mathcal{B}}_n[(u - x_0)^2] \\
 &\quad + \frac{1}{2} (f_{xy}(x_0, y_0) + f_{yx}(x_0, y_0)) \tilde{\mathcal{B}}_n[(u - x_0)(v - y_0)] \\
 &\quad + \frac{1}{2} f_{yy}(x_0, y_0) \tilde{\mathcal{B}}_n[(v - y_0)^2] + \tilde{\mathcal{B}}_n \left[ \|(u, v) - (x_0, y_0)\|^2 h(u, v) \right],
 \end{aligned}$$

where we have omitted  $\Omega$  for brevity. We deal with each term separately.

From Lemma 5.1 (ii), we get  $\tilde{\mathcal{B}}_n[u - x_0]|_{u=x_0} = 0$ . Next, from the proof of Lemma 5.1 (iii), we have

$$\tilde{\mathcal{B}}_n[v - y_0] = \tilde{B}_n[\phi_2(x) - \phi_1(x), I] \frac{y - \phi_1(x)}{\phi_2(x) - \phi_1(x)} + \tilde{B}_n[\phi_1(x), I] - y_0.$$

But using (5.3), we get

$$\begin{aligned}
 \tilde{\mathcal{B}}_n[v - y_0]|_{(u,v)=(x_0,y_0)} &= \left( \phi_2(x_0) - \phi_1(x_0) + \mathcal{O}\left(\frac{1}{n}\right) \right) \frac{y_0 - \phi_1(x_0)}{\phi_2(x_0) - \phi_1(x_0)} \\
 &\quad + \phi_1(x_0) + \mathcal{O}\left(\frac{1}{n}\right) - y_0 = \mathcal{O}\left(\frac{1}{n}\right).
 \end{aligned}$$

Similarly,

$$\begin{aligned}
 \tilde{\mathcal{B}}_n[(u - x_0)^2]|_{(u,v)=(x_0,y_0)} &= \mathcal{O}\left(\frac{1}{n}\right), \\
 \tilde{\mathcal{B}}_n[(u - x_0)(v - y_0)]|_{(u,v)=(x_0,y_0)} &= \mathcal{O}\left(\frac{1}{n}\right), \\
 \tilde{\mathcal{B}}_n[(v - y_0)^2]|_{(u,v)=(x_0,y_0)} &= \mathcal{O}\left(\frac{1}{n}\right).
 \end{aligned}$$

Now we deal with the last term

$$\begin{aligned} & \tilde{\mathcal{B}}_n \left[ \|(u, v) - (x_0, y_0)\|^2 h(u, v) \right] \\ &= \sum_{k=0}^n \sum_{j=0}^{n_k} \tilde{F} \left( \frac{k}{n}, \frac{j}{n_k} \right) \tilde{H} \left( \frac{k}{n}, \frac{j}{n_k} \right) \tilde{p}_{n,k}(x; I) \tilde{p}_{n_k,j}(y; [\phi_1, \phi_2]), \end{aligned}$$

where

$$\begin{aligned} & \tilde{F} \left( \frac{k}{n}, \frac{j}{n_k} \right) \\ &= \left( (b-a) \frac{k}{n} + a - x_0 \right)^2 + \left( \left( \tilde{\phi}_2 \left( \frac{k}{n} \right) - \tilde{\phi}_1 \left( \frac{k}{n} \right) \right) \frac{j}{n_k} + \tilde{\phi}_1 \left( \frac{k}{n} \right) - y_0 \right)^2, \end{aligned}$$

and

$$\tilde{H} \left( \frac{k}{n}, \frac{j}{n_k} \right) = h \left( (b-a) \frac{k}{n} + a, \left( \tilde{\phi}_2 \left( \frac{k}{n} \right) - \tilde{\phi}_1 \left( \frac{k}{n} \right) \right) \frac{j}{n_k} + \tilde{\phi}_1 \left( \frac{k}{n} \right) \right).$$

Fix a real number  $\varepsilon > 0$ . Then, there is a real number  $\delta > 0$  such that if  $\|(u, v) - (x_0, y_0)\| < \delta$ , then  $|h(u, v)| < \varepsilon$ . Let  $S_\delta$  be the set of  $k$  and  $j$  such that  $\frac{1}{\delta^2} \tilde{F} \left( \frac{k}{n}, \frac{j}{n_k} \right) > 1$ . Then,

$$\begin{aligned} & \sum_{(k,j) \in S_\delta} \tilde{p}_{n,k}(x_0; I) \tilde{p}_{n_k,j}(y_0; [\phi_1, \phi_2]) \\ &< \frac{1}{\delta^2} \sum_{(k,j) \in S_\delta} \tilde{F} \left( \frac{k}{n}, \frac{j}{n_k} \right) \tilde{p}_{n,k}(x_0; I) \tilde{p}_{n_k,j}(y_0; [\phi_1, \phi_2]) \\ &\leq \frac{1}{\delta^2} \left( \tilde{\mathcal{B}}_n[(u-x_0)^2] + \tilde{\mathcal{B}}_n[(v-y_0)^2] \right) \Big|_{(u,v)=(x_0,y_0)} \\ &= \mathcal{O} \left( \frac{1}{n} \right). \end{aligned}$$

Moreover, we have

$$\begin{aligned} & \sum_{(k,j) \notin S_\delta} \tilde{F} \left( \frac{k}{n}, \frac{j}{n_k} \right) \left| \tilde{H} \left( \frac{k}{n}, \frac{j}{n_k} \right) \right| \tilde{p}_{n,k}(x; I) \tilde{p}_{n_k,j}(y; [\phi_1, \phi_2]) \\ &< \varepsilon \sum_{(k,j) \notin S_\delta} \tilde{F} \left( \frac{k}{n}, \frac{j}{n_k} \right) \tilde{p}_{n,k}(x_0; I) \tilde{p}_{n_k,j}(y_0; [\phi_1, \phi_2]) \leq \mathcal{O} \left( \frac{1}{n} \right). \end{aligned}$$

Thus,

$$\begin{aligned} & \left| \tilde{\mathcal{B}}_n \left[ \left\| (u, v) - (x_0, y_0) \right\|^2 h(u, v) \right] \right| \\ & \leq \sum_{(k, j) \in S_\delta} \left| \tilde{F} \left( \frac{k}{n}, \frac{j}{n_k} \right) \tilde{H} \left( \frac{k}{n}, \frac{j}{n_k} \right) \right| \tilde{p}_{n,k}(x; I) \tilde{p}_{n_k, j}(y; [\phi_1, \phi_2]) \\ & \quad + \sum_{(k, j) \notin S_\delta} \left| \tilde{F} \left( \frac{k}{n}, \frac{j}{n_k} \right) \tilde{H} \left( \frac{k}{n}, \frac{j}{n_k} \right) \right| \tilde{p}_{n,k}(x; I) \tilde{p}_{n_k, j}(y; [\phi_1, \phi_2]) \\ & \leq M \sum_{(k, j) \in S_\delta} \tilde{p}_{n,k}(x_0; I) \tilde{p}_{n_k, j}(y_0; [\phi_1, \phi_2]) + \mathcal{O} \left( \frac{1}{n} \right) \\ & \leq \mathcal{O} \left( \frac{1}{n} \right), \end{aligned}$$

where

$$M = \sup_{(u, v) \in \Omega} \left\| \left\| (u, v) - (x_0, y_0) \right\|^2 h(u, v) \right\|.$$

Putting all the above together, we get

$$\left| \tilde{\mathcal{B}}_n [f(x, y), \Omega] \Big|_{(x, y) = (x_0, y_0)} - f(x_0, y_0) \right| \leq \mathcal{O} \left( \frac{1}{n} \right),$$

and the result follows. □

### 6 Shifted Bernstein-Type Operators

We define the *shifted* bivariate Bernstein-type operator. Let  $\phi_1$  and  $\phi_2$  be two continuous functions, and let  $I = [a, b]$  be an interval such that  $\phi_1 < \phi_2$  on  $I$ . Let  $\Omega \subset \mathbb{R}^2$  be the domain bounded by the curves  $y = \phi_1(x)$ ,  $y = \phi_2(x)$ , and the straight lines  $x = a$ ,  $x = b$ . Let

$$T(x, y) = (\tau(x), \sigma_x(y)), \quad (x, y) \in \Omega,$$

where  $\tau$  is any continuously differentiable function on  $I$ , such that  $\tau(a) = a$ ,  $\tau(b) = b$ , and  $\tau'(x) > 0$  for  $x \in I$ , and for each fixed  $x \in I$ ,  $\sigma_x$  is any continuously differentiable function on  $[\phi_1(x), \phi_2(x)]$ , such that  $\sigma_x(\phi_1(x)) = \phi_1(x)$ ,  $\sigma_x(\phi_2(x)) = \phi_2(x)$ , and  $\sigma'_x(y) > 0$  for  $y \in [\phi_1(x), \phi_2(x)]$ .

For every function  $f(x, y)$  defined on  $\Omega$ , define the function

$$\tilde{F}^T(u, v; \Omega) = f \circ T^{-1} \left( (b - a)u + a, (\tilde{\phi}_2(u) - \tilde{\phi}_1(u))v + \tilde{\phi}_1(u) \right),$$

for  $0 \leq u \leq 1$ , and  $0 \leq v \leq 1$ , where  $\tilde{\phi}_i, i = 1, 2$ , are defined in (5.1).

The shifted bivariate Bernstein-type operator is defined as

$$\tilde{\mathcal{E}}_n^T [f(x, y), \Omega] = \sum_{k=0}^n \sum_{j=0}^{n_k} \tilde{F}^T \left( \frac{k}{n}, \frac{j}{n_k}; \Omega \right) \tilde{p}_{n,k}(\tau(x); I) \tilde{p}_{n_k,j}(\sigma_x(y); [\phi_1, \phi_2]),$$

for  $(x, y) \in \Omega$ , where  $n_k = n - k$  or  $n_k = k$  for  $0 \leq k \leq n$ .

Written in terms of the univariate classical Bernstein basis, we get

$$\begin{aligned} \tilde{\mathcal{E}}_n^T [f(x, y), \Omega] &= \sum_{k=0}^n \sum_{j=0}^{n_k} \tilde{F}^T \left( \frac{k}{n}, \frac{j}{n_k}; \Omega \right) p_{n,k} \left( \frac{\tau(x) - a}{b - a} \right) \\ & p_{n_k,j} \left( \frac{\sigma_x(y) - \phi_1(x)}{\phi_2(x) - \phi_1(x)} \right). \end{aligned}$$

**Proposition 6.1** For every function  $f(x, y)$  defined on  $\Omega$ ,

$$\lim_{n \rightarrow +\infty} \tilde{\mathcal{E}}_n^T [f(x, y), \Omega] = f(x, y).$$

**Proof** Let  $u = \tau(x)$  and, for each  $x \in I, v = \sigma_x(y)$ . Then,

$$\begin{aligned} \tilde{\mathcal{E}}_n^T [f(x, y), \Omega] &= \sum_{k=0}^n \sum_{j=0}^{n_k} \tilde{F} \left( \frac{k}{n}, \frac{j}{n_k}; a, b \right) \tilde{p}_{n,k}(u; I) \tilde{p}_{n_k,j}(v; [\phi_1, \phi_2]) \\ &= \tilde{\mathcal{B}}_n [(f \circ T)(u, v), \Omega]. \end{aligned}$$

From Theorem 5.3, we have  $\tilde{\mathcal{B}}_n [(f \circ T)(u, v), \Omega] = \tilde{\mathcal{B}}_n [f(x, y), \Omega]$  converges uniformly to  $f(x, y)$ . Hence,  $\tilde{\mathcal{E}}_n^T [f(x, y), \Omega]$  converges uniformly to  $f(x, y)$ .  $\square$

Now, we study shifted Bernstein-type operators defined on each quadrant of  $\mathbf{B}^2$ , denoted by  $B_i$  for  $i = 1, 2, 3, 4$ . We will choose  $T$  and  $n_k$  such that, for any function, the approximation given by Bernstein-type operators on each quadrant is a polynomial.

(i) For  $x \in [0, 1]$ , let  $\tau(x) = x^2$  and, for each fixed value of  $x$ , let  $\sigma_x(y) = y^2/\sqrt{1-x^2}$ . Let  $n_k = n - k, \phi_1(x) = 0$ , and  $\phi_2(x) = \sqrt{1-x^2}$ . Then,

$$\begin{aligned} \tilde{p}_{n,k}(x^2; [0, 1]) &= \binom{n}{k} x^{2k} (1-x^2)^{n-k}, \\ \tilde{p}_{n-k,j}(\sigma_x(y); [\phi_1, \phi_2]) &= \frac{1}{(1-x^2)^{n-k}} \binom{n-k}{j} y^{2j} (1-x^2-y^2)^{n-k-j}, \\ \tilde{F}^T(u, v; B_1) &= f \left( \sqrt{u}, \sqrt{(1-u)v} \right), \end{aligned}$$

where  $B_1 = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 \leq 1, x, y \geq 0\}$ . Then,

$$\tilde{\mathcal{E}}_n^T [f(x, y), B_1] = \sum_{k=0}^n \sum_{j=0}^{n-k} f \left( \sqrt{\frac{k}{n}}, \sqrt{\frac{j}{n}} \right) \binom{n}{k} x^{2k} y^{2j} (1-x^2-y^2)^{n-k-j}.$$

(ii) For  $x \in [-1, 0]$ , let  $\tau(x) = -x^2$  and, for each fixed value of  $x$ , let  $\sigma_x(y) = y^2/\sqrt{1-x^2}$ . Let  $n_k = k$ ,  $\phi_1(x) = 0$ , and  $\phi_2(x) = \sqrt{1-x^2}$ . Then,

$$\begin{aligned} \tilde{p}_{n,k}(-x^2; [-1, 0]) &= \binom{n}{k} (1-x^2)^k x^{2n-2k}, \\ \tilde{p}_{k,j}(\sigma_x(y); [\phi_1, \phi_2]) &= \frac{1}{(1-x^2)^k} \binom{k}{j} y^{2j} (1-x^2-y^2)^{k-j}, \\ \tilde{F}^T(u, v; B_2) &= f(-\sqrt{1-u}, \sqrt{uv}), \end{aligned}$$

where  $B_2 = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 \leq 1, x \leq 0, y \geq 0\}$ . Then,

$$\begin{aligned} \tilde{\mathcal{C}}_n^T[f(x, y), B_2] &= \sum_{k=0}^n \sum_{j=0}^k f\left(-\sqrt{1-\frac{k}{n}}, \sqrt{\frac{j}{n}}\right) \\ &\quad \binom{n}{k} \binom{k}{j} x^{2n-2k} y^{2j} (1-x^2-y^2)^{k-j}. \end{aligned}$$

(iii) For  $x \in [-1, 0]$ , let  $\tau(x) = -x^2$  and, for each fixed value of  $x$ , let  $\sigma_x(y) = -y^2/\sqrt{1-x^2}$ . Let  $n_k = k$ ,  $\phi_1(x) = -\sqrt{1-x^2}$ , and  $\phi_2(x) = 0$ . Then,

$$\begin{aligned} \tilde{p}_{n,k}(-x^2; [-1, 0]) &= \binom{n}{k} (1-x^2)^k x^{2n-2k}, \\ \tilde{p}_{k,j}(\sigma_x(y); [\phi_1, \phi_2]) &= \frac{1}{(1-x^2)^k} \binom{k}{j} (1-x^2-y^2)^j y^{2k-2j}, \\ \tilde{F}^T(u, v; B_3) &= f\left(-\sqrt{1-u}, -\sqrt{u(1-v)}\right), \end{aligned}$$

where  $B_3 = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 \leq 1, x, y \leq 0\}$ . Then,

$$\begin{aligned} \tilde{\mathcal{C}}_n^T[f(x, y), B_3] &= \sum_{k=0}^n \sum_{j=0}^k f\left(-\sqrt{1-\frac{k}{n}}, -\sqrt{\frac{k-j}{n}}\right) \\ &\quad \binom{n}{k} \binom{k}{j} x^{2n-2k} y^{2k-2j} (1-x^2-y^2)^j. \end{aligned}$$

(iv) For  $x \in [0, 1]$ , let  $\tau(x) = x^2$  and, for each fixed value of  $x$ , let  $\sigma_x(y) = -y^2/\sqrt{1-x^2}$ . Let  $n_k = n-k$ ,  $\phi_1(x) = -\sqrt{1-x^2}$ , and  $\phi_2(x) = 0$ . Then,

$$\begin{aligned} \tilde{p}_{n,k}(x^2; [0, 1]) &= \binom{n}{k} x^{2k} (1-x^2)^{n-k}, \\ \tilde{p}_{n-k,j}(\sigma_x(y); [\phi_1, \phi_2]) &= \frac{1}{(1-x^2)^{n-k}} \binom{n-k}{j} (1-x^2-y^2)^j y^{2n-2k-2j}, \\ \tilde{F}^T(u, v; B_4) &= f\left(\sqrt{u}, -\sqrt{(1-u)(1-v)}\right), \end{aligned}$$



where  $B_4 = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 \leq 1, x \geq 0, y \leq 0\}$ . Then,

$$\begin{aligned} \tilde{\mathcal{C}}_n^T[f(x, y), B_4] &= \sum_{k=0}^n \sum_{j=0}^{n-k} f\left(\sqrt{\frac{k}{n}}, -\sqrt{1 - \frac{k+j}{n}}\right) \\ &\quad \binom{n}{k \ j} x^{2k} (1 - x^2 - y^2)^j y^{2n-2k-2j}. \end{aligned}$$

Similar to (4.1), we can define a piecewise Bernstein-type operator on  $\mathbf{B}^2$  as follows:

$$\bar{\mathcal{C}}_n[f(x, y), \mathbf{B}^2] = \begin{cases} \tilde{\mathcal{C}}_n^T[f(x, y), B_1], & (x, y) \in B_1, \\ \tilde{\mathcal{C}}_n^T[f(x, y), B_2], & (x, y) \in B_2, \\ \tilde{\mathcal{C}}_n^T[f(x, y), B_3], & (x, y) \in B_3, \\ \tilde{\mathcal{C}}_n^T[f(x, y), B_4], & (x, y) \in B_4. \end{cases} \tag{6.1}$$

The proof of the following proposition is similar to that of Proposition 4.1.

**Proposition 6.2** *For any function  $f$  on  $\mathbf{B}^2$ ,  $\bar{\mathcal{C}}_n[f(x, y), \mathbf{B}^2]$  is a continuous function on  $\mathbf{B}^2$ .*

### 7 Better Order of Approximation

In [3], Butzer studied a certain linear combination of univariate Bernstein operators that, under certain conditions, give a better order of approximation compared with the classical operators. For a bounded function  $f(x)$  defined on  $[0, 1]$ , Butzer considers the linear combination of Bernstein polynomials

$$\mathfrak{L}_n^{[2k]} f(x) = \alpha_k B_{2k_n} f(x) + \alpha_{k-1} B_{2^{k-1}_n} f(x) + \dots + \alpha_0 B_n f(x), \quad k \geq 0,$$

where the constants  $\alpha_j = \alpha_j(k)$  satisfy

$$\alpha_k + \alpha_{k-1} + \dots + \alpha_0 = 1.$$

The polynomials  $\mathfrak{L}_n^{[2k]} f(x)$  satisfy the recurrence relation

$$\begin{aligned} (2^k - 1)\mathfrak{L}_n^{[2k]} f(x) &= 2^k \mathfrak{L}_{2n}^{[2k-2]} f(x) - \mathfrak{L}_n^{[2k-2]} f(x), \\ \mathfrak{L}_n^{[0]} f(x) &= B_n f(x), \end{aligned} \tag{7.1}$$

and, if  $f^{(2k)}$  exists at a point  $x \in [0, 1]$ , then

$$\mathfrak{L}_n^{[2k-2]} f(x) - f(x) = \mathcal{O}\left(n^{-k}\right), \quad k \geq 0.$$

Using (7.1), we can obtain the following explicit expressions for the constants  $\alpha_j$ 's,

$$\alpha_j = \prod_{\substack{i=0 \\ i \neq j}}^k \frac{2^j}{2^j - 2^i}, \quad 0 \leq j \leq k. \tag{7.2}$$

In [14], May considers a slightly more general operator  $S_n^{[k]}[f(x); d_0, d_1, \dots, d_k] \equiv S_n^{[k]}[f(x); d_j]$  given by

$$\begin{aligned} S_n^{[k]}[f(x); d_j] &= \sum_{j=0}^k \prod_{\substack{i=0 \\ i \neq j}}^k \frac{d_j}{d_j - d_i} B_{d_j n} f(x), \\ S_n^{[0]}[f(x); d_j] &= B_{d_0 n} f(x), \end{aligned} \tag{7.3}$$

where  $d_0, d_1, \dots, d_k$  are positive integers. Notice that if  $d_j = 2^j$ , then  $S_n^{[k]}[f(x); d_j] = \mathfrak{L}_n^{[2^k]} f(x)$ . Moreover, if  $d_j = j + 1$ , then  $S_n^{[2^k-1]}[f(x); j + 1]$  is a polynomial of degree  $2^k n$ . However, May proved that if  $f^{(2^{k+1})}$  exists, then  $S_n^{[2^k-1]}[f(x), j + 1]$  converges to  $f(x)$  at a rate of  $n^{-2^k}$  and, hence,  $S_n^{[2^k-1]}[f(x), j + 1]$  and  $\mathfrak{L}_n^{[2^k]} f(x)$  are polynomials of the same degree, but  $S_n^{[2^k-1]}[f(x), j + 1]$  has a faster rate of convergence than  $\mathfrak{L}_n^{[2^k]} f(x)$ . Observe that for  $k = 1$ , we have  $S_n^{[2^k-1]}[f(x); j + 1] = S_n^{[k]}[f(x); 2^j]$ .

Motivated by the construction in (7.3), we define the following bivariate operators

$$\begin{aligned} \tilde{S}_n^{[k]}[f(x, y); d_j] &= \sum_{j=0}^k \prod_{\substack{i=0 \\ i \neq j}}^k \frac{d_j}{d_j - d_i} \tilde{\mathcal{B}}_{d_j n}[f(x, y), \mathbf{B}^2], \\ \tilde{S}_n^{[0]}[f(x, y); d_j] &:= \tilde{\mathcal{B}}_{d_0 n}[f(x, y), \mathbf{B}^2], \end{aligned} \tag{7.4}$$

and

$$\begin{aligned} \tilde{R}_n^{[k]}[f(x, y), d_j] &:= \sum_{j=0}^k \prod_{\substack{i=0 \\ i \neq j}}^k \frac{d_j}{d_j - d_i} \overline{\mathcal{C}}_{d_j n}[f(x, y), \mathbf{B}^2], \\ \tilde{R}_n^{[0]}[f(x, y); d_j] &:= \overline{\mathcal{C}}_{d_0 n}[f(x, y), \mathbf{B}^2]. \end{aligned} \tag{7.5}$$

Although we do not study the approximation behavior of these operators here, the numerical experiments in the following section suggest a better rate of convergence than  $\tilde{\mathcal{B}}_n$  and  $\overline{\mathcal{C}}_n$ .

## 8 Numerical Experiments

In this section, we present numerical experiments where we compare the shifted Bernstein–Stancu operator  $\tilde{\mathcal{B}}_n$  on  $\mathbf{B}^2$ , and the shifted Bernstein-type operator  $\overline{\mathcal{C}}_n$  in (6.1). To do this, we consider different functions defined on  $\mathbf{B}^2$ . For each function  $f(x, y)$ , we compute  $\tilde{\mathcal{B}}_n[f(x, y), \mathbf{B}^2]$  and  $\overline{\mathcal{C}}_n[f(x, y), \mathbf{B}^2]$ . We use a set of points randomly distributed on the unit disk (generated by *mesh* function in Mathematica) to compare the function to its approximations. For  $\tilde{\mathcal{B}}_n[f(x, y), \mathbf{B}^2]$ , we use 630 points  $(x_i, y_i)$ . We set  $z_i = f(x_i, y_i)$ ,  $1 \leq i \leq 630$ , and  $\hat{z}_i$  equal to the value of  $\tilde{\mathcal{B}}_n[f(x, y), \mathbf{B}^2]$  at the respective point  $(x_i, y_i)$ , and compute the root-mean-square error (RMSE) as follows:

$$\text{RMSE}(f, \tilde{\mathcal{B}}_n) = \sqrt{\sum_{i=1}^{630} \frac{(z_i - \hat{z}_i)^2}{630}}.$$

Similarly, for  $\overline{\mathcal{C}}_n[f(x, y), \mathbf{B}^2]$ , we use randomly distributed 1082 points  $(\bar{x}_j, \bar{y}_j)$ . We set  $w_j = f(\bar{x}_j, \bar{y}_j)$ ,  $1 \leq j \leq 1082$ , and  $\bar{w}_j$  equal to the value of  $\overline{\mathcal{C}}_n[f(x, y), \mathbf{B}^2]$  at the respective point  $(\bar{x}_j, \bar{y}_j)$ , and compute the RMSE as follows:

$$\text{RMSE}(f, \overline{\mathcal{C}}_n) = \sqrt{\sum_{j=1}^{1082} \frac{(w_j - \bar{w}_j)^2}{1082}}.$$

In each case, we plot the RMSE for increasing values of  $n$  using Mathematica.

For each operator, the set of points used to compute the RMSE consists of a fixed number of points. On the other hand, the number of mesh points used to represent each operator depends on  $n$ .

We represent  $\overline{\mathcal{C}}_n[f(x, y), \mathbf{B}^2]$  on each quadrant using different colors as shown in Fig. 4. We take  $n = 100$ , then the mesh required to obtain the operator for each quadrant consists of 20200 points.

For  $\tilde{\mathcal{B}}_n[f(x, y), \mathbf{B}^2]$ , we take  $n = 200$ . Then, the mesh required to obtain the operator for all the unit disk consists of 40401 points.

We note that the operator  $\overline{\mathcal{C}}_n$  requires two evaluations at the mesh points on the common boundaries of two adjacent quadrants. Therefore, the operator  $\tilde{\mathcal{B}}_n$  needs a smaller number of evaluations than the operator  $\overline{\mathcal{C}}_n$  since, for a fixed  $n$ ,  $\tilde{\mathcal{B}}_n$  and  $\overline{\mathcal{C}}_n$  are composed of  $(n + 1)^2$  and  $2(n + 1)(n + 2)$  evaluations, respectively.

Additionally, we compute the RMSE for  $\tilde{\mathcal{S}}_n^{[1]}[f(x, y), 2^j]$  and  $\tilde{\mathcal{R}}_n^{[1]}[f(x, y), 2^j]$  using the same set of randomly distributed points as before.

We note that, from the definition, we have

$$\begin{aligned}\tilde{\mathcal{S}}_n^{[1]}[f(x, y); 2^j] &= 2\tilde{\mathcal{B}}_{2n}[f(x, y), \mathbf{B}^2] - \tilde{\mathcal{B}}_n[f(x, y), \mathbf{B}^2], \\ \tilde{\mathcal{R}}_n^{[1]}[f(x, y), 2^j] &= 2\overline{\mathcal{C}}_{2n}[f(x, y), \mathbf{B}^2] - \overline{\mathcal{C}}_n[f(x, y), \mathbf{B}^2].\end{aligned}$$





Fig. 4 Left: Mesh for  $\overline{\mathcal{C}}_n$  with  $n = 15$ . Color code for disk quadrants ( $B_1$  red;  $B_2$  green;  $B_3$  yellow;  $B_4$  purple). Right: Mesh for  $\widetilde{\mathcal{B}}_n$  with  $n = 15$

Fig. 5 Graph of  $f(x, y) = x \sin(5x - 6y) + y$  on  $\mathbf{B}^2$




For a fixed  $n$ , the mesh required to obtain  $\widetilde{\mathcal{B}}_{2n}$  (respectively,  $\overline{\mathcal{C}}_{2n}$ ) is a refinement of the mesh required for  $\widetilde{\mathcal{B}}_n$  (respectively,  $\overline{\mathcal{C}}_n$ ). Therefore,  $\widetilde{\mathcal{R}}_n^{[1]}[f(x, y), 2^j]$  and  $\widetilde{\mathcal{S}}_n^{[1]}[f(x, y), 2^j]$  require  $(2n + 1)^2$  and  $2(2n + 1)(2n + 2)$  evaluations, respectively.

### 8.1 Example 1

First, we consider the continuous function

$$f(x, y) = x \sin(5x - 6y) + y, \quad (x, y) \in \mathbf{B}^2.$$

The graph of  $f(x, y)$  is shown in Fig. 5, and the approximations  $\overline{\mathcal{C}}_n[f(x, y), \mathbf{B}^2]$  and  $\mathcal{B}_n[f(x, y), \mathbf{B}^2]$  are shown in Fig. 6. We list the RMSE of both approximations for different values of  $n$  in Table 1 and plot them together in Fig. 7, where the characteristic slow convergence inherited from the univariate Bernstein operators is observed. Moreover, the corresponding RMSEs are shown in Table 2 and Fig. 8 for  $\widetilde{\mathcal{S}}_n^{[1]}[f(x, y), 2^j]$  and  $\widetilde{\mathcal{R}}_n^{[1]}[f(x, y), 2^j]$ , where a seemingly better approximation behavior can be observed.




**Fig. 6** Approximations overlapped with the graph of  $f(x, y)$ . Left:  $\overline{\mathcal{E}}_n[f(x, y), \mathbf{B}^2]$ . Right:  $\widetilde{\mathcal{B}}_n[f(x, y), \mathbf{B}^2]$

**Table 1** RMSE for different values of  $n$

$n$	$\overline{\mathcal{E}}_n[f(x, y), \mathbf{B}^2]$	$\widetilde{\mathcal{B}}_n[f(x, y), \mathbf{B}^2]$
10	0.191411	0.30623
20	0.117881	0.209091
30	0.0860663	0.16182
40	0.0682511	0.132416
50	0.0568288	0.112151
60	0.0488602	0.0972969
70	0.0429694	0.0859318
80	0.0384267	0.0769527

**Fig. 7** Plot of RMSE in Table 1



### 8.2 Example 2

Now, we consider the continuous periodic function


$$g(x, y) = \sin(10x + y), \quad (x, y) \in \mathbf{B}^2.$$

Its graph is shown in Fig. 9. It can be observed in Fig. 10 that the approximation error for both operators is larger at the maximum and minimum values of the function. Table

**Table 2** RMSE for different values of  $n$

$n$	$\tilde{\mathcal{R}}_n^{[1]}[f(x, y); 2^j]$	$\tilde{\mathcal{S}}_n^{[1]}[f(x, y); 2^j]$
10	0.0592126	0.14241
20	0.0306588	0.0666483
30	0.0217412	0.0389884
40	0.0171251	0.0258265
50	0.014188	0.0185806
60	0.0121117	0.0141754
70	0.0105426	0.0112949
80	0.00929964	0.009302

**Fig. 8** Plot of RMSE in Table 2






**Table 3** RMSE for different values of  $n$

$n$	$\tilde{\mathcal{C}}_n[g(x, y), \mathbf{B}^2]$	$\tilde{\mathcal{B}}_n[g(x, y), \mathbf{B}^2]$
10	0.535344	0.700146
20	0.366915	0.613427
30	0.278477	0.526227
40	0.225091	0.454904
50	0.189454	0.398559
60	0.163967	0.353775
70	0.144812	0.317628
80	0.129872	0.287968


3 and Fig. 11 contain further evidence of this larger error. Moreover, in comparison with the previous example, it seems that the rate convergence of  $\tilde{\mathcal{C}}_n[g(x, y), \mathbf{B}^2]$  is significantly faster than the rate of convergence of  $\tilde{\mathcal{B}}_n[g(x, y), \mathbf{B}^2]$ . Table 4 and Fig. 12 show the errors corresponding to  $\tilde{\mathcal{R}}_n^{[1]}[f(x, y), 2^j]$  and  $\tilde{\mathcal{S}}_n^{[1]}[f(x, y), 2^j]$ . In comparison with  $\tilde{\mathcal{B}}_n$  and  $\tilde{\mathcal{C}}_n$ ,  $\tilde{\mathcal{R}}_n^{[1]}$ , and  $\tilde{\mathcal{S}}_n^{[1]}$  appear to have a better approximation behavior.

**Fig. 9** Graph of  $g(x, y) = \sin(10x + y)$  on  $\mathbf{B}^2$



**Fig. 10** Approximations overlapped with the graph of  $g(x, y)$ . Left:  $\overline{\mathcal{C}}_n[g(x, y), \mathbf{B}^2]$ . Right:  $\tilde{\mathcal{B}}_n[g(x, y), \mathbf{B}^2]$


**Fig. 11** Plot of RMSE in Table 3



**Table 4** RMSE for different values of  $n$

$n$	$\tilde{\mathcal{R}}_n^{[1]}[g(x, y); 2^j]$	$\tilde{\mathcal{S}}_n^{[1]}[g(x, y); 2^j]$
10	0.231422	0.53511
20	0.116893	0.300211
30	0.0799078	0.18369
40	0.061706	0.122657
50	0.0506578	0.0873721
60	0.043083	0.0652876
70	0.0374756	0.0505948
80	0.0331074	0.0403428

**Fig. 12** Plot of RMSE in Table 4



### 8.3 Example 3



Here, we consider the continuous function

$$h(x, y) = e^{x^2-y^2} - xy, \quad (x, y) \in \mathbf{B}^2,$$

(see Fig. 13). Both approximations are shown in Fig. 14, and their respective RMSEs are listed in Table 5 and plotted in Fig. 15. Observe that, in this case, the RMSEs for both approximations are significantly smaller than in the previous examples. Moreover, based on Fig. 15, it seems that for sufficiently large values of  $n$ , the rate of convergence of both approximations is considerably similar to each other. Table 6 and Fig. 16 also show similar approximation behavior between  $\tilde{\mathcal{S}}_n^{[1]}[f(x, y), 2^j]$  and  $\tilde{\mathcal{R}}_n^{[1]}[f(x, y), 2^j]$ .



**Fig. 13** Graph of  $h(x, y) = e^{x^2-y^2} - xy$



**Fig. 14** Approximations overlapped with the graph of  $h(x, y)$ . Left:  $\overline{\mathcal{C}}_n[h(x, y), \mathbf{B}^2]$ . Right:  $\widetilde{\mathcal{B}}_n[h(x, y), \mathbf{B}^2]$

**Table 5** RMSE for different values of  $n$

$n$	$\overline{\mathcal{C}}_n[h(x, y), \mathbf{B}^2]$	$\widetilde{\mathcal{B}}_n[h(x, y), \mathbf{B}^2]$
10	0.0505862	0.140837
20	0.0293585	0.0685387
30	0.0213945	0.0455634
40	0.017105	0.0342737
50	0.0143844	0.0275514
60	0.0124871	0.0230843
70	0.0110789	0.0198962
80	0.00998647	0.0175041

### 8.4 Example 4

In this numerical example, we are interested in observing the behavior of shifted Bernstein-type and shifted Bernstein–Stancu operators at jump discontinuities.

Let us consider the following discontinuous function:

$$\eta(x, y) = \begin{cases} 1, & \text{if } x^2 + y^2 < 0.5, \\ 0, & \text{if } 0.5 \leq x^2 + y^2 \leq 0.8, \\ 0.5, & \text{if } 0.8 < x^2 + y^2 \leq 1. \end{cases}$$

The graph of  $\eta(x, y)$  is shown in Fig. 17 and the approximations are shown in Fig. 14. It is interesting to observe the behavior of the approximations at the points of jump

Fig. 15 Plot of RMSE in Table 5




Table 6 RMSE for different values of n

n	$\tilde{\mathcal{R}}_n^{[1]}[h(x, y); 2^j]$	$\tilde{\mathcal{S}}_n^{[1]}[h(x, y); 2^j]$
10	0.015677	0.0188267
20	0.00931191	0.0087729
30	0.0068375	0.00610222
40	0.00545586	0.00479186
50	0.00455264	0.00399096
60	0.00390752	0.00343958
70	0.00341937	0.0030301
80	0.00303472	0.00270966

Fig. 16 Plot of RMSE in Table 6




Fig. 17 Graph of  $\eta(x, y)$





Fig. 18 Approximations overlapped with the graph of  $\eta(x, y)$ . Left:  $\overline{\mathcal{C}}_n[\eta(x, y), \mathbf{B}^2]$ . Right:  $\widetilde{\mathcal{B}}_n[\eta(x, y), \mathbf{B}^2]$





Fig. 19 Cross-sectional view of the approximations for increasing values of  $n$ . Left:  $\overline{\mathcal{C}}_n[\eta(x, y), \mathbf{B}^2]$ . Right:  $\mathcal{B}_n[\eta(x, y), \mathbf{B}^2]$

discontinuities and, thus, we have included Fig. 19, where we show a cross-sectional view of the approximations with increasing values of  $n$ . As in the univariate case, it seems that the Gibbs phenomenon does not occur. Finally, Table 7 and Fig. 20 expose a significantly slow convergence rate for this discontinuous function in comparison with the previous continuous examples. As can be seen in Table 8 and Fig. 21, it seems

**Table 7** RMSE for different values of  $n$

$n$	$\overline{\mathcal{E}}_n[\eta(x, y), \mathbf{B}^2]$	$\widetilde{\mathcal{B}}_n[\eta(x, y), \mathbf{B}^2]$
10	0.216588	0.270366
20	0.175754	0.243468
30	0.156563	0.223305
40	0.144805	0.210916
50	0.136559	0.205949
60	0.130305	0.192988
70	0.125319	0.193887
80	0.121205	0.187675


**Fig. 20** Plot of RMSE in Table 7



**Table 8** RMSE for different values of  $n$

$n$	$\widehat{\mathcal{R}}_n^{[1]}[\eta(x, y); 2^j]$	$\widetilde{\mathcal{S}}_n^{[1]}[\eta(x, y); 2^j]$
10	0.173582	0.203794
20	0.132669	0.149787
30	0.121201	0.140103
40	0.108429	0.122498
50	0.102915	0.120874
60	0.0959724	0.111816
70	0.0923721	0.111529
80	0.0878217	0.105521

Fig. 21 Plot of RMSE in Table 8



that a better approximation can be obtained with the operators  $\tilde{\mathcal{S}}_n^{[1]}[f(x, y), 2^j]$  and  $\tilde{\mathcal{R}}_n^{[1]}[f(x, y), 2^j]$ .

**Acknowledgements** The authors are grateful to the reviewer for their valuable comments and suggestions which led us to improve this work.

**Funding** Open Access funding provided thanks to the CRUE-CSIC agreement with Springer Nature.

## Declarations

**Conflict of interest** The authors declare no conflict of interests.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

- Berens, H., Schmid, H.J., Xu, Y.: Bernstein–Durrmeyer polynomials on a simplex. *J. Approx. Theory* **68**(3), 247–261 (1992)
- Bernstein, S.: Demonstration di Théorème de Weierstrass fondée sur le calcul des probabilités. *Commun. Soc. Math. Kharkov* (2), 13 (1912–1913), 1, 2
- Butzer, P.: Linear combinations of Bernstein polynomials. *Can. J. Math.* **5**, 559–567 (1953)
- Cárdenas-Morales, D., Garrancho, P., Muñoz-Delgado, F.J.: Shape preserving approximation by Bernstein-type operators which fix polynomials. *Appl. Math. Comput.* **182**, 1615–1622 (2006)
- Cárdenas-Morales, D., Garrancho, P., Raşa, I.: Bernstein-type operators that preserve polynomials. *Comput. Math. with Appl.* **62**, 158–163 (2011)
- Derriennic, M.M.: Sur l'approximation de fonctions intégrables sur  $[0, 1]$  par des polynômes de Bernstein modifiés. *J. Approx. Theory* **31**(4), 325–343 (1981)

7. Derriennic, M.M.: On multivariate approximation by Bernstein-type polynomials. *J. Approx. Theory* **45**(2), 155–166 (1985)
8. Dunkl, C.F., Xu, Y.: *Orthogonal Polynomials of Several Variables*. Encyclopedia of Mathematics and its Applications, vol. 155, 2nd edn. Cambridge University Press, Cambridge (2014)
9. Durrmeyer, J. L.: Une formule d'inversion de la transformée de Laplace. Applications à la théorie des moments, Thèse 3e cycle, Fac. des Sciences de l'Université de Paris (1967)
10. Gonska, H., Pițul, P., Rașa, I.: General King-type operators. *Results Math.* **53**(3–4), 279–286 (2009)
11. Kantorovitch, L. V.: Sur certains développements suivants les polynômes de la forme de S. Bernstein, I, II, C. R. Acad. Sci. USSR, 563–568, 595–600 (1930)
12. King, J.P.: Positive linear operators which preserve  $x^2$ . *Acta Math. Hungar.* **99**(3), 203–208 (2003)
13. Lorentz, G.G.: *Bernstein Polynomials*. Chelsea Publishing Company, New York (1997)
14. May, C.: Saturation and inverse theorems for combinations of a class of exponential-type operators. *Can. J. Math.* **28**(6), 1224–1250 (1976)
15. Sablonnière, P.: Opérateurs de Bernstein–Jacobi et polynômes orthogonaux, Publ. ANO 37, Laboratoire de Calcul, Université de Lille (1981)
16. Schurer, F.: On the approximation of functions of many variables with linear positive operators. *Nederl. Akad. Wetensch. Proc. Ser. A 66 = Indag. Math.* **25**, 313–327 (1963)
17. Stancu, D.D.: A method for obtaining polynomials of Bernstein type of two variables. *Am. Math. Mon.* **70**(3), 260–264 (1963)
18. Volkov, V.I.: On the convergence of sequences of linear positive operators in the space of continuous functions of two variables. *Dokl. Akad. Nauk. SSSR (NS)* **115**, 17–19 (1957)

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.